

Anti-Rhp51 / Rad51 (*S. pombe*) antibody, rabbit polyclonal, validated

63-001 100 µl

Validation: Specificity has been validated by western blotting with rhp51 deletion mutant (Fig.1)

Storage temmperature: Shipped at 4°C and store at -20°C

Reactivity: Schizosaccharomyces pombe

Immunogen: Purified recombinant full-length Rhp51 protein

Application

1. Western blotting (1/ 2,000~1/5,000) Fig.1
2. Immunoprecipitation (1/100-1/500)
3. Chromatin Immuno-Precipitation (Assay dependent)
4. Immunofluorescence staining (1/500 dilution). Fig. 2

Form: Rabbit antiserum added with 0.09 % sodium azide

Background: Rhp51 protein of *Schizosaccharomyces pombe* (fission yeast) is a functional and structural homolog of *E.coli* RecA protein and Rad51 proteins of eukaryotes, which play a major role in genetic recombination and recombination repair by mediating strand exchange reaction between homologous DNA strands.

Data Link UniProtKB/Swiss-Prot [P36601](#) (RAD51_SCHPO)

References :This product has been used in the following publication

1. Akamatsu Y et al. Two different Swi5-containing protein complexes are involved in mating-type switching and recombination repair in fission yeast. [Proc Natl Acad Sci U S A.](#) 2003 Dec 23;100(26):15770-5. **WB, IP (*S. pombe*)**
2. Kibe T et al. Fission yeast Rhp51 is required for the maintenance of telomere structure in the absence of the Ku heterodimer. [Nucleic Acids Res.](#) 2003 Sep 1;31(17):5054-63. **ChIP (*S. pombe*)**
- 2.Lambert S et al "Gross chromosomal rearrangements and elevated recombination at an inducible site-specific replication fork barrier" [Cell](#) 121: 689-702 (2005) PMID: [15935756](#) **IF (*S. pombe*)**
3. Morishita T et al "Role of the Schizosaccharomyces pombe F-Box DNA helicase in processing recombination intermediates" [Mol Cell Biol](#) 25: 8074-8083 (2005) PMID: [16135799](#) **IF (*S.pombe*)**
4. Haruta N et al "The Swi5-Sfr1 complex stimulates Rhp51/Rad51-and Dmc1-mediated DNA strand exchange in vitro" [Nat Struc Mol Biol](#) 13: 823-830 (2006) PMID: [16921379](#) **WB, IP (*S. pombe*)**
5. Akamatsu Y et al. Fission yeast Swi5/Sfr1 and Rhp55/Rhp57 differentially regulate Rhp51-dependent recombination outcomes. [EMBO J.](#) 2007 Mar 7;26(5):1352-62. **IF (*S.***

pombe)

6. Polakova S et al. Dbl2 Regulates Rad51 and DNA Joint Molecule Metabolism to Ensure Proper Meiotic Chromosome Segregation. *PLoS Genet.* 2016 Jun 15;12(6):e1006102. IF (*S. pombe*)




Fig.1 Western blot analysis of Rhp51 in the whole cell extracts. M: Molecular size markers (kD) Lane 1: Wild-type strain Lane2: Rhp51 deletion mutant strain




Fig. 2 Rhp51 foci formation observed after DNA damage: *S. pombe* cells without or with MMS (0.025%) treatment for 1 h were processed for indirect immunofluorescence staining with anti-Rhp51 antibody (1/500 dilution).