

#### PRODUCT DATA SHEET

# Anti-Rabbit IgG (H+L) - 60nm Gold Conjugate (1ml, OD10)

Catalog No. AC-60-01-15 Size: 1ml

### **Description**

Affinity isolated anti-rabbit antibody produced in goat and coupled to 60nm gold nanoparticle. Suitable for use in immunoblotting, lateral flow, light microscopy, and electron microscopy applications procedures for secondary detection of rabbit antibody labeled samples.

Provides a permanent and sensitive label (100pg or less) when used in conjunction with Cytodiagnostics membrane and microscopy silver enhancer kits, see related product below.

Concentration: 0.5 mg/ml (OD=10)

Conjugated Antibody: Goat affinity purified anti-rabbit


IgG(H+L)

Clonality: Polyclonal

Storage Buffer: 20mM Tris (pH 8.0), 150mM NaCl,

20% glycerol (v/v), 1% BSA **Working Dilution:** 1:30 – 1:300





## **Storage**

Store undiluted in storage buffer at 2-8° C. Stable for at least 4 months if stored as specified.

## **Product Safety and Handling**

This product is for R&D use only, not for drug, household, or other uses. Please review the material safety datasheet (MSDS) available online for proper safety and handling procedures.

#### **Related Products**

Silver Enhancer Kit for Membranes Cat No. SR-01-02 Silver Enhancer Kit for Microscopy Cat No. SR-01-01

## Standard Immunogold Dot-Blot Protocol

(Adapted from Moeremans et al. [1])

- Spot one microlitre drops of a serial dilution of your protein (100-0.1ng) in PBS supplemented with 50ug/ml of BSA on nitrocellulose or PVDF membrane.
- 2. Let protein drops dry into the membrane.
- Block Membrane for 30 minutes using 1% (w/v) dry milk in 1X PBS at room temperature.
- 4. Incubate with primary antibody for 2 hours at room temperature.
- 5. Wash membrane 3x5 minutes with blocking solution prepared as above.
- Incubate for 2 hours (or longer for increased sensitivity) with secondary gold conjugate diluted 1:10 (OD=0.3) times with blocking solution (0.2% Blocking Solution). See Tech Note #102 for preparation of a gold conjugate.
- 7. Wash 3x5 minutes as above.
- 8. Dry membrane and record data.
- 9. (OPTIONAL) Proceed with <u>silver</u> enhancement to improve sensitivity.






Figure 1. Example dot-blot assay for Cytodiagnostics streptavidin gold conjugate (top left) and our streptavidin silver conjugate (top right) before and after enhancement using Cytodiagnostics silver enhancement kit for membranes. Bottom picture illustrates and highlights the difference in appearance (color) of 50nm anti-human IgG noble metal nanoparticle conjugates prepared using NHS-activated gold nanoparticles, NHS-activated gold nanoparticles, and NHS-activated silver nanoparticles,

#### References

respectively.

1. M. Moeremans, et al., Journal of Immunological Methods, 1984, 74, 353

## **Ordering Information**

For ordering call 866-344-3954 or visit us online at www.cytodiagnostics.com