

Langford Locks Kidlington Oxford OX5 1LH United Kingdom

Tel: +44 (0)1865 595230

Antibody Datasheet

Product Name: Rabbit anti *Borrelia burgdorferi* CRASP-2

Product: Purified rabbit anti CRASP-2 antibody, unconjugated

Product Type: Polyclonal

Isotype: Rabbit IgG

Product code: PAB21448-100

Batch Number: R001218

Amount: 0.1 ml (1.0 mg/mL by UV absorbance at 280 nm)

Physical State: Lyophilized

Buffer: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2

Reconstitution Vol: 100 μL

Reconstitution Buffer: Deionized water (or equivalent)

Preservative: 0.01% (w/v) Sodium Azide

Immunogen: Recombinant MBP tagged B. burgdorferi CRASP-2 protein

Purification: Protein-A purified and cross-adsorbed against MBP from monospecific

antiserum by chromatography

Specificity: This antibody is specific for *Borrelia burgdorferi* CRASP-2 protein. A BLAST

analysis was used to suggest reactivity with CRASP-2 from *B. burgdorferi* sources based on 100% homology with the immunizing sequence. Partial cross-reactivity is expected against *B. garinii*, *B. spielmanii*, and *valaisiana* sources based on 91-89% homology. Cross-reactivity with CRASP-2 from

other sources has not been determined.






Applications: ELISA (1:1000), WB (1:1000)

Storage: Store at 4° C prior to restoration. For extended storage aliquot contents and

freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. Antibody is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to

immediate use.



Western Blot showing detection of 0.1 μg of recombinant CRASP-2 protein. Lane 1: Molecular weight markers. Lane 2: MBP-CRASP-2 fusion protein (arrow; expected MW = 67.8 kDa). Lane 3: MBP alone. Protein was run on a 4-20% gel, transferred to 0.45 nitrocellulose. After blocking with 1% BSA-TTBS overnight at 4°C, primary antibody was used at 1:1000 at room temperature for 30 min. HRPconjugated Goat-Anti-Rabbit secondary antibody was used at 1:40,000 in blocking buffer and imaged on the VersaDoc™ MP 4000 imaging system (Bio-Rad).

