

Langford Locks Kidlington Oxford OX5 1LH United Kingdom Tel: +44 (0)1865 595230

Antibody Datasheet

Product Name:	Rabbit anti Borrelia burgdorferi DbpA
Product:	Purified rabbit anti DbpA antibody, unconjugated
Product Type:	Polyclonal
lsotype:	Rabbit IgG
Product code:	PAB21449-100
Batch Number:	R001218
Amount:	0.1 ml (1.0 mg/mL by UV absorbance at 280 nm)
Physical State:	Lyophilized
Buffer:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Reconstitution Vol:	100 μL
Reconstitution Buffer:	Deionized water (or equivalent)
Preservative:	0.01% (w/v) Sodium Azide
Immunogen:	Recombinant MBP tagged <i>B. burgdorferi</i> DbpA protein
Purification:	Protein-A purified and cross-adsorbed against MBP from monospecific antiserum by chromatography
Specificity:	This antibody is specific for Borrelia burgdorferi DbpA protein. A BLAST analysis was used to suggest cross-reactivity with DbpA from Borrelia burgdorferi sources based on 100% homology with the immunizing sequence. Partial reactivity is expected against Borrelia garinii sources based on 60-80% homology. Cross-reactivity with DbpA from other sources has not been determined.






Applications:

ELISA (1:5000), WB (1:1000)

Storage:

Store at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. Antibody is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.



Western blot showing detection of 0.1 µg recombinant proteins in western blot. Lane 1: Molecular weight markers. Lane 2: MBP-DbpA fusion protein (arrow; expected MW: 60.9 kDa). Lane 3: DbpA, MBP removed by TEV cleavage. Lane 4: MBP alone. Protein was run on a 4-20% gel, then transferred to 0.45 µm nitrocellulose. After blocking with 1% BSA-TTBS (overnight at 4°C), primary antibody used 1:1000 was at at room temperature for 30 min. HRPconjugated Goat-Anti-Rabbit secondary antibody was used at 1:40,000 in MB-070 blocking buffer and imaged on the VersaDoc[™] MP 4000 imaging system (Bio-Rad).

