

Langford Locks
Kidlington
Oxford
OX5 1LH
United Kingdom
Tel: +44 (0)1865 595230

Antibody Datasheet

Product Name: Rabbit anti Borrelia burgdorferi P39

Product: Purified rabbit anti P39 antibody, unconjugated

Product Type: Polyclonal

Isotype: Rabbit IgG

Product code: PAB21457-25

Batch Number: R001218

Amount: 25 μl (1.0 mg/mL by UV absorbance at 280 nm)

Physical State: Liquid (sterile filtered)

Buffer: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2

Preservative: 0.01% (w/v) Sodium Azide

Immunogen: Recombinant MBP tagged B. burgdorferi P39 protein

Purification: Protein-A purified and cross-adsorbed against MBP from monospecific

antiserum by chromatography

Specificity: This antibody is specific for *Borrelia burgdorferi* P39 protein. A BLAST analysis

was used to suggest cross-reactivity with p39 from B. burgdorferi and B. afzelii sources based on 100% homology with the immunizing sequence. Partial reactivity is expected against *B. garinii* sources based on 88-95% homology and to *B. spielmanii* A14S based on 86-96% homology. Cross-

reactivity with p39 from other sources has not been determined.


Applications: ELISA (1:5000), WB (1:1000)





Storage:

Store vial at -20° C or below prior to opening. To minimize loss of volume, dilute 1:10 by adding 225 μ L of the buffer stated above directly to the vial. Recap, mix thoroughly and briefly centrifuge to collect the volume at the bottom of the vial. Use this intermediate dilution when calculating final dilutions as recommended above. Store the vial at -20°C or below after dilution. Avoid cycles of freezing and thawing.



Western blot showing detection of 0.1 μg of recombinant P39 protein (expected size ~35.4 kDa in size). Lane 1: Molecular weight markers. Lane 2: MBP-P39 fusion protein (expected MW: 77.8 kDa). Lane 3: MBP alone. Protein was run on a 4-20% gel, then transferred to 0.45 µm nitrocellulose. After blocking with 1% BSA-TTBS overnight at 4°C, primary antibody was used at 1:1000 at room temperature for 30 min. HRP-conjugated Goat-Anti-Rabbit secondary antibody was used at 1:40,000 in blocking buffer and imaged on the VersaDoc™ MP 4000 imaging system (Bio-Rad).