

219 North Arkansas Street Rogers, AR 72756 Phone: (479) 636-4361; 800-643-3426 Fax: (479) 636-3562 Email: <u>biosales@pel-freez.com</u> Website: <u>www.pelfreez-bio.com</u>

PRODUCT DATA SHEET

Product Name: ANTI-PHOSPHO-Ser^{62,67} SYNAPSIN I ANTIBODY

Product Code:

P40029-100 100 μL

Pack Size:

Description: Synapsin I plays a key role in synaptic plasticity in brain (Feng et al., 2002; Nayak et al., 1996). This effect is due in large part to the ability of the synapsins to regulate the availability of synaptic vesicles for release. The role of synapsin in synaptic plasticity and in synaptogensis is regulated by phosphorylation (Jovanovic et al., 2001; Kao et al., 2002). Ser⁵⁴⁹ along with Ser⁶² and Ser⁶⁷ are the sites of Synapsin I that are phosphorylated by MAP kinase (Czernik et al., 1987; Jovanovic et al., 1996).

Physical State:	Liquid; Buffer contents: 10 mM HEPES (pH 7.5), 150 mM NaCl, 100 µg per mL BSA and 50% glycerol
Storage/Stability:	Stable at -20 °C for at least 1 year. For long term storage -20 °C is recommended
Purification Method:	Prepared from rabbit serum by affinity purification via sequential chromatography on phospho- and dephosphopeptide affinity columns.
Shipping Conditions:	Domestic: Blue Ice International: Blue Ice or Dry Ice

Host Species:	Rabbit	(Polyclonal)
<i>M</i> r (kDa):	78	

Immunogen: Phosphopeptide corresponding to amino acid residues surrounding the phospho-Ser^{62,67} of synapsin I. Specific for ~78k synapsin I doublet phosphorylated at Ser^{62,67}. Immunolabeling of the synapsin I band is blocked by preadsorption with the phospho-peptide used as antigen but not be the corresponding dephospho-peptide.

Species Reactivity: The antibody has been directly tested for reactivity in Western blots with rat tissue. It is anticipated that the antibody will react with mouse, and bovine tissues based on the fact that these species have 100% homology with the amino acid sequence used as antigen.

Recommended Antibody Dilutions:

WB: 1:1000

References:

- 1) Czernik AJ et al. (1987) *Proc Natl Acad Sci (USA)* 84:7518-7522.
- 2) Feng J et al. (2002) J Neurosci 22:4372-4380.
- 3) Jovanovic JN et al. (1996) *Neurobiology* 93:3679-3683.
- 4) Jovanovic JN et al. (2001) *J Neurosci* 21:7944-7953.
- 5) Kao HT et al. (2002) Nature Neurosci 5:431-437.
- 6) Nayak AS et al. (1996) *Proc Natl Acad Sci (USA)* 93:15451-15456.

Western Blot

Rat cortex lysate showing specific labeling of the ~78k synapsin protein phosphorylated at Ser^{62,67} (Control). Immunolabeling is blocked by preadsorption with the phospho-peptide used as antigen (Peptide) but not by the corresponding dephosphopeptide (not shown).



Application Key: WB – Western Blot IF – Immunofluorescence IHC – Immunohistochemistry IP - Immunoprecipitation

P/N: 74111 Rev 01

Page 1 of 1

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES.