

## Bassoon

Cat.No. 141-0P; control protein, 100 µg protein (lyophilized)

### Data Sheet

Reconstitution/ Storage	100 µg protein, lyophilized. For reconstitution add 100 µl H <sub>2</sub> O to get a 1mg/ml solution in TBS. Then aliquot and store at -20°C until use.
Immunogen	Recombinant protein corresponding to AA 3608 to 3938 from rat Bassoon (UniProt Id: O88778)
Recommended dilution	Optimal concentrations should be determined by the end-user.
matching antibodies	141 002, 141 003, 141 011, 141 021
Remarks	This control protein consists of the recombinant protein that has been used for immunization. It has been tested in preadsorption experiments and blocks efficiently and specifically the corresponding signal in Western blots. The amount of protein needed for efficient blocking depends on the titer and on the affinity of the antibody to the antigen.

**TO BE USED IN VITRO / FOR RESEARCH ONLY**  
**NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS**

**Bassoon** is a large protein which consists of an N-terminal Zn<sup>2+</sup> finger and several piccolo-bassoon homology domains (PBH-domains). It is generally found together with piccolo, a related huge multi-domain protein of the CAZ (cytoskeletal matrix assembled at active zones). Bassoon was suggested to be a scaffolding element of the presynapse but deletion experiments in mice have shown that bassoon is also involved in synaptic vesicle cycling. Probably bassoon interacts with other protein factors via its Zn<sup>2+</sup> domain but the potential partners have not been determined yet.

### Selected General References

Functional regions of the presynaptic cytomatrix protein bassoon: significance for synaptic targeting and cytomatrix anchoring. Dresbach T, Hempelmann A, Spilker C, tom Dieck S, Altmann WD, Zuschratter W, Garner CC, Gundelfinger ED. *Molecular and cellular neurosciences* (2003) 23(2): 279-91.

Unitary assembly of presynaptic active zones from Piccolo-Bassoon transport vesicles. Shapira M, Zhai RG, Dresbach T, Bresler T, Torres VI, Gundelfinger ED, Ziv NE, Garner CC. *Neuron* (2003) 38(2): 237-52.

Functional inactivation of a fraction of excitatory synapses in mice deficient for the active zone protein bassoon. Altmann WD, tom Dieck S, Sokolov M, Meyer AC, Sigler A, Brakebusch C, Fässler R, Richter K, Boeckers TM, Potschka H, Brandt C, et al. *Neuron* (2003) 37(5): 787-800.

The presynaptic active zone protein bassoon is essential for photoreceptor ribbon synapse formation in the retina. Dick O, tom Dieck S, Altmann WD, Ammermüller J, Weiler R, Garner CC, Gundelfinger ED, Brandstätter JH. *Neuron* (2003) 37(5): 775-86.

Localization of the presynaptic cytomatrix protein Piccolo at ribbon and conventional synapses in the rat retina: comparison with Bassoon.

Dick O, Hack I, Altmann WD, Garner CC, Gundelfinger ED, Brandstätter JH. *The Journal of comparative neurology* (2001) 439(2): 224-34.

Membrane association of presynaptic cytomatrix protein bassoon.

Sanmartí-Vila L, tom Dieck S, Richter K, Altmann W, Zhang L, Volkhardt W, Zimmermann H, Garner CC, Gundelfinger ED, Dresbach T. *Biochemical and biophysical research communications* (2000) 275(1): 43-6.

Bassoon, a novel zinc-finger CAG/glutamine-repeat protein selectively localized at the active zone of presynaptic nerve terminals.

tom Dieck S, Sanmartí-Vila L, Langnaese K, Richter K, Kindler S, Soyke A, Wex H, Smalla KH, Kämpf U, Fränzer JT, Stumm M, et al. *The Journal of cell biology* (1998) 142(2): 499-509.