

Bassoon

Cat.No. 141 016; Polyclonal chicken antibody, 200 µl antibody (lyophilized)

Data Sheet

Reconstitution/ Storage	200 µl antibody, lyophilized. For reconstitution add 200 µl H ₂ O, then aliquot and store at -20°C until use.
Applications	WB: not recommended IP: not tested yet ICC: 1 : 500 IHC: 1 : 500 IHC-P/FFPE: 1 : 500
Immunogen	Recombinant protein corresponding to AA 756 to 1001 from rat Bassoon (UniProt Id: O88778)
Reactivity	Reacts with: rat (O88778), mouse (O88737). Other species not tested yet.
Specificity	Specific for bassoon.

TO BE USED IN VITRO / FOR RESEARCH ONLY
NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS

Bassoon is a large protein which consists of an N-terminal Zn²⁺ finger and several piccolo-bassoon homology domains (PBH-domains). It is generally found together with piccolo, a related huge multi-domain protein of the CAZ (cytoskeletal matrix assembled at active zones).

Bassoon was suggested to be a scaffolding element of the presynapse but deletion experiments in mice have shown that bassoon is also involved in synaptic vesicle cycling. Probably bassoon interacts with other protein factors via its Zn²⁺ domain but the potential partners have not been determined yet.

Selected References SYSY Antibodies

Coherent Activity between the Prelimbic and Auditory Cortex in the Slow-Gamma Band Underlies Fear Discrimination.
Concina G, Cambiaghi M, Renna A, Sacchetti B
The Journal of neuroscience : the official journal of the Society for Neuroscience (2018) 38(39): 8313-8328. **IHC; tested species: rat**

Selected General References

Functional regions of the presynaptic cytomatrix protein bassoon: significance for synaptic targeting and cytomatrix anchoring.
Dresbach T, Hempelmann A, Spilker C, tom Dieck S, Altmann WD, Zuschratter W, Garner CC, Gundelfinger ED
Molecular and cellular neurosciences (2003) 23(2): 279-91.

Unitary assembly of presynaptic active zones from Piccolo-Bassoon transport vesicles.
Shapira M, Zhai RG, Dresbach T, Bresler T, Torres VI, Gundelfinger ED, Ziv NE, Garner CC
Neuron (2003) 38(2): 237-52.

Functional inactivation of a fraction of excitatory synapses in mice deficient for the active zone protein bassoon.
Altmann WD, tom Dieck S, Sokolov M, Meyer AC, Sigler A, Brakebusch C, Fässler R, Richter K, Boeckers TM, Potschka H, Brandt C, et al.
Neuron (2003) 37(5): 787-800.

The presynaptic active zone protein bassoon is essential for photoreceptor ribbon synapse formation in the retina.
Dick O, tom Dieck S, Altmann WD, Ammermüller J, Weiler R, Garner CC, Gundelfinger ED, Brandstätter JH
Neuron (2003) 37(5): 775-86.

Localization of the presynaptic cytomatrix protein Piccolo at ribbon and conventional synapses in the rat retina: comparison with Bassoon.

Dick O, Hack I, Altmann WD, Garner CC, Gundelfinger ED, Brandstätter JH
The Journal of comparative neurology (2001) 439(2): 224-34.

Membrane association of presynaptic cytomatrix protein bassoon.
Sanmartí-Vila L, tom Dieck S, Richter K, Altmann W, Zhang L, Volknandt W, Zimmermann H, Garner CC, Gundelfinger ED, Dresbach T
Biochemical and biophysical research communications (2000) 275(1): 43-6.

Bassoon, a novel zinc-finger CAG/glutamine-repeat protein selectively localized at the active zone of presynaptic nerve terminals.
tom Dieck S, Sanmartí-Vila L, Langnaese K, Richter K, Kindler S, Soyke A, Wex H, Smalla KH, Kämpf U, Fränzer JT, Stumm M, et al.
The Journal of cell biology (1998) 142(2): 499-509.