

Calbindin D28k

Rudolf-Wissell-Str. 28
37079 Göttingen, Germany
Phone: +49 551-50556-0
Fax: +49 551-50556-384
E-mail: sales@sysy.com
Web: www.sysy.com

Cat.No. 214 004; Polyclonal Guinea pig antibody, 100 µl antiserum (lyophilized)

Data Sheet

Reconstitution/ Storage	100 µl antiserum, lyophilized. For reconstitution add 100 µl H ₂ O, then aliquot and store at -20°C until use.
Applications	WB: 1 : 1000 up to 1 : 5000 (AP staining) IP: yes ICC: 1 : 500 IHC: 1 : 200 up to 1 : 500 IHC-P/FFPE: 1 : 200
Immunogen	Recombinant protein corresponding to AA 3 to 251 from human CalbindinD28k (UniProt Id: P05937)
Reactivity	Reacts with: human (P05937), rat (P07171), mouse (P12658), monkey, ape, cow. Other species not tested yet.
Specificity	Specific for calbindin D28k.
matching control	214-0P

TO BE USED IN VITRO / FOR RESEARCH ONLY
NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS

Two isoforms of the vitamin D-dependent Ca-binding proteins have been described so far: **calbindin D28k**, also referred to as CALB 1, D-28k, and CAB 27, and calbindin D29k, also known as calretinin. These proteins are expressed in cells that have to handle a high calcium influx such as brain, bone, teeth, inner ear and others. Calbindins are believed to regulate cellular activity by suppressing or buffering intracellular calcium

Selected References SYSY Antibodies

- Cell age-specific vulnerability of neurons to anesthetic toxicity.
Hofacer RD, Deng M, Ward CG, Joseph B, Hughes EA, Jiang C, Danzer SC, Loepke AW
Annals of neurology (2013) 73(6): 695-704. **IHC**
- Non-canonical heterogeneous cellular distribution and co-localization of CaMKIIα and CaMKIIβ in the spinal superficial dorsal horn.
Larsson M
Brain structure & function (2018) 223(3): 1437-1457. **IHC; tested species: rat**
- Hypothalamic CNTF volume transmission shapes cortical noradrenergic excitability upon acute stress.
Alpár A, Zahola P, Hanics J, Hevesi Z, Korchyńska S, Benevento M, Pifl C, Zachar G, Perugini J, Severi I, Leitgeb P, et al.
The EMBO journal (2018) : . **IHC; tested species: mouse**
- Abolished perineuronal nets and altered parvalbumin-immunoreactivity in the nucleus reticularis thalami of wildtype and 3xTg mice after experimental stroke.
Härtig W, Appel S, Suttkus A, Grosche J, Michalski D
Neuroscience (2016) 337: 66-87. **IHC**
- Development of early-born γ-Aminobutyric acid hub neurons in mouse hippocampus from embryogenesis to adulthood.
Villette V, Guigue P, Picardo MA, Sousa VH, Leprince E, Lachamp P, Malvache A, Tressard T, Cossart R, Baude A
The Journal of comparative neurology (2016) 524(12): 2440-61. **IHC; tested species: mouse**
- An excitatory GABA loop operating in vivo.
Astorga G, Bao J, Marty A, Augustine GJ, Franconville R, Jalil A, Bradley J, Llano I
Frontiers in cellular neuroscience (2015) 9: 275. **IHC**
- Functional and Physical Interaction of Diacylglycerol Kinase ζ with Protein Kinase Ca Is Required for Cerebellar Long-Term Depression.
Lee D, Yamamoto Y, Kim E, Tanaka-Yamamoto K
The Journal of neuroscience : the official journal of the Society for Neuroscience (2015) 35(46): 15453-65. **IHC**
- Raf kinase inhibitory protein is required for cerebellar long-term synaptic depression by mediating PKC-dependent MAPK activation.
Yamamoto Y, Lee D, Kim Y, Lee B, Seo C, Kawasaki H, Kuroda S, Tanaka-Yamamoto K
The Journal of neuroscience : the official journal of the Society for Neuroscience (2012) 32(41): 14254-64. **IHC**
- Postmitotic neurons develop a p21-dependent senescence-like phenotype driven by a DNA damage response.
Jurk D, Wang C, Miwa S, Maddick M, Korolchuk V, Tsolou A, Gonos ES, Thrasivoulou C, Saffrey MJ, Cameron K, von Zglinicki T, et al.
Aging cell (2012) 11(6): 996-1004. **IHC**
- Spontaneous activity promotes synapse formation in a cell-type-dependent manner in the developing retina.
Soto F, Ma X, Cecil JL, Vo BQ, Culican SM, Kerschensteiner D
The Journal of neuroscience : the official journal of the Society for Neuroscience (2012) 32(16): 5426-39. **IHC**
- Slow age-dependent decline of doublecortin expression and BrdU labeling in the forebrain from lesser hedgehog tenrecs.
Alpár A, Künzle H, Gärtner U, Popkova Y, Bauer U, Grosche J, Reichenbach A, Härtig W
Brain research (2010) 1330: 9-19. **IHC**
- ### Selected General References
- Influence of the "open field" exposure on calbindin D28K, calretinin, and parvalbumin containing cells in the rat midbrain - developmental study.
Klejbor I, Ludkiewicz B, Domaradzka-Pytel B, Spodnik JH, Dziwiatkowski J, Moryś J
Journal of physiology and pharmacology : an official journal of the Polish Physiological Society (2006) 57(1): 149-64.
- Calbindin D-28 and microtubule-associated protein-2: their use as sensitive immunohistochemical markers of cerebellar neurotoxicity in a regulatory toxicity study.
Haworth R, McCormack N, Selway S, Pilling AM, Williams TC
Experimental and toxicologic pathology : official journal of the Gesellschaft für Toxikologische Pathologie (2006) 57(5-6): 419-26.
- Mutational analysis of dendritic Ca²⁺ kinetics in rodent Purkinje cells: role of parvalbumin and calbindin D28k.
Schmidt H, Stiefel KM, Racay P, Schwaller B, Eilers J
The Journal of physiology (2003) 551(Pt 1): 13-32.
- Calbindin in cerebellar Purkinje cells is a critical determinant of the precision of motor coordination.
Barski JJ, Hartmann J, Rose CR, Hoebeek F, Mörl K, Noll-Hussong M, De Zeeuw CI, Konnerth A, Meyer M
The Journal of neuroscience : the official journal of the Society for Neuroscience (2003) 23(8): 3469-77.