

## SynCAM 2

Cat.No. 243 203; Polyclonal rabbit antibody, 50 µg specific antibody (lyophilized)

### Data Sheet

Reconstitution/ Storage	50 µg specific antibody, lyophilized. Affinity purified with the immunogen. Rabbit serum albumin was added for stabilization. For reconstitution add 50 µl H <sub>2</sub> O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C until use.
Applications	<b>WB:</b> 1 : 1000 (AP staining) <b>IP:</b> not tested yet <b>ICC:</b> 1 : 100 <b>IHC:</b> 1 : 100 <b>IHC-P/FFPE:</b> not tested yet
Immunogen	Synthetic peptide corresponding to AA 167 to 181 from mouse SynCAM2 (UniProt Id: Q8BLQ9)
Reactivity	Reacts with: rat (Q1WIM2), mouse (Q8BLQ9). Other species not tested yet.
Specificity	Specific for synCAM 2. (K.O. verified)
matching control	243-2P

**TO BE USED IN VITRO / FOR RESEARCH ONLY**  
**NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS**

Development of synapses involves transsynaptic interactions of dedicated synaptic adhesion molecules like neuroligins, neurexins and synCAMs (**S**ynaptic cell **a**dhesion **m**olecules). Four SynCAM isoforms have been described so far. All share a common domain structure and contain three extracellular Ig domains, a single transmembrane region, and a short COOH-terminal cytoplasmic tail. **SynCAM 2** has been shown to be associated to myelinated axons and **synCAM 3** can serve as a receptor for herpes viruses.

### Selected References SYSY Antibodies

Cadm2 regulates body weight and energy homeostasis in mice.  
Yan X, Wang Z, Schmidt V, Gauert A, Willnow TE, Heinig M, Poy MN  
Molecular metabolism (2018) 8: 180-188. **WB, ICC; KO verified; tested species: mouse**

### Selected General References

Aberrations of a cell adhesion molecule CADM4 in renal clear cell carcinoma.  
Nagata M, Sakurai-Yageta M, Yamada D, Goto A, Ito A, Fukuhara H, Kume H, Morikawa T, Fukayama M, Homma Y, Murakami Y, et al.  
International journal of cancer (2012) 130(6): 1329-37.

Localization of Cadm2a and Cadm3 proteins during development of the zebrafish nervous system.  
Hunter PR, Nikolaou N, Odermatt B, Williams PR, Drescher U, Meyer MP  
The Journal of comparative neurology (2011) 519(11): 2252-70.

The cell adhesion nectin-like molecules (Nect1) 1 and 4 suppress the growth and tumorigenic ability of colon cancer cells.  
Raveh S, Gavert N, Spiegel I, Ben-Ze'ev A  
Journal of cellular biochemistry (2009) 108(1): 326-36.

The adhesion molecule Nect1-3/SynCAM-2 localizes to myelinated axons, binds to oligodendrocytes and promotes cell adhesion.  
Pellissier F, Gerber A, Bauer C, Ballivet M, Ossipov V  
BMC neuroscience (2007) 8: 90.

Nectin-like proteins mediate axon Schwann cell interactions along the internode and are essential for myelination.  
Maurel P, Einheber S, Galinska J, Thaker P, Lam I, Rubin MB, Scherer SS, Murakami Y, Gutmann DH, Salzer JL  
The Journal of cell biology (2007) 178(5): 861-74.

SynCAMs organize synapses through heterophilic adhesion.  
Fogel AI, Akins MR, Krupp AJ, Stagi M, Stein V, Biederer T  
The Journal of neuroscience : the official journal of the Society for Neuroscience (2007) 27(46): 12516-30.

Crystal structure of the V domain of human Nectin-like molecule-1/Syncam3/Tsll1/lgsf4b, a neural tissue-specific immunoglobulin-like cell-cell adhesion molecule.  
Dong X, Xu F, Gong Y, Gao J, Lin P, Chen T, Peng Y, Qiang B, Yuan J, Peng X, Rao Z, et al.  
The Journal of biological chemistry (2006) 281(15): 10610-7.

Nectin-like molecule-1/TSLL1/SynCAM3: a neural tissue-specific immunoglobulin-like cell-cell adhesion molecule localizing at non-junctional contact sites of presynaptic nerve terminals, axons and glia cell processes.  
Kakunaga S, Ikeda W, Itoh S, Deguchi-Tawarada M, Ohtsuka T, Mizoguchi A, Takai Y  
Journal of cell science (2005) 118(Pt 6): 1267-77.

SynCAM, a synaptic adhesion molecule that drives synapse assembly.  
Biederer T, Sara Y, Mozhayeva M, Atasoy D, Liu X, Kavalali ET, Südhof TC  
Science (New York, N.Y.) (2002) 297(5586): 1525-31.