



Rudolf-Wissell-Str. 28
37079 Göttingen, Germany
Phone: +49 551-50556-0
Fax: +49 551-50556-384
E-mail: sales@sysy.com
Web: www.sysy.com

SynCAM 3

Cat.No. 243 303; Polyclonal rabbit antibody, 50 µg specific antibody (lyophilized)

Data Sheet

| | |
|-------------------------|---|
| Reconstitution/ Storage | 50 µg specific antibody, lyophilized. Affinity purified with the immunogen. Rabbit serum albumin was added for stabilization. For reconstitution add 50 µl H ₂ O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C until use. |
| Applications | WB: 1 : 1000 (AP staining) IP: not tested yet ICC: not tested yet IHC: not tested yet IHC-P/FFPE: not tested yet |
| Immunogen | Synthetic peptide corresponding to AA 171 to 185 from mouse SynCAM3 (UniProt Id: Q99N28) |
| Reactivity | Reacts with: rat (Q1WIM3), mouse (Q99N28). Other species not tested yet. |
| Specificity | Specific for synCAM 3. |
| matching control | 243-3P |

Selected General References

- Aberrations of a cell adhesion molecule CADM4 in renal clear cell carcinoma.
Nagata M, Sakurai-Yageta M, Yamada D, Goto A, Ito A, Fukuhara H, Kume H, Morikawa T, Fukayama M, Homma Y, Murakami Y, et al.
International journal of cancer (2012) 130(6): 1329-37.
- Localization of Cadm2a and Cadm3 proteins during development of the zebrafish nervous system.
Hunter PR, Nikolau N, Odermatt B, Williams PR, Drescher U, Meyer MP
The Journal of comparative neurology (2011) 519(11): 2252-70.
- The cell adhesion nectin-like molecules (Necl) 1 and 4 suppress the growth and tumorigenic ability of colon cancer cells.
Raveh S, Gavert N, Spiegel I, Ben-Ze'ev A
Journal of cellular biochemistry (2009) 108(1): 326-36.
- The adhesion molecule Necl-3/SynCAM-2 localizes to myelinated axons, binds to oligodendrocytes and promotes cell adhesion.
Pellissier F, Gerber A, Bauer C, Ballivet M, Ossipow V
BMC neuroscience (2007) 8: 90.
- Nectin-like proteins mediate axon Schwann cell interactions along the internode and are essential for myelination.
Maurel P, Einheber S, Galinska J, Thaker P, Lam I, Rubin MB, Scherer SS, Murakami Y, Gutmann DH, Salzer JL
The Journal of cell biology (2007) 178(5): 861-74.
- SynCAMs organize synapses through heterophilic adhesion.
Fogel Al, Akins MR, Krupp AJ, Stagi M, Stein V, Biederer T
The Journal of neuroscience : the official journal of the Society for Neuroscience (2007) 27(46): 12516-30.
- Crystal structure of the V domain of human Nectin-like molecule-1/Syncam3/Tsl1/Igsf4b, a neural tissue-specific immunoglobulin-like cell-cell adhesion molecule.
Dong X, Xu F, Gong Y, Gao J, Lin P, Chen T, Peng Y, Qiang B, Yuan J, Peng X, Rao Z, et al.
The Journal of biological chemistry (2006) 281(15): 10610-7.
- Nectin-like molecule-1/TSLL1/SynCAM3: a neural tissue-specific immunoglobulin-like cell-cell adhesion molecule localizing at non-junctional contact sites of presynaptic nerve terminals, axons and glia cell processes.
Kakunaga S, Ikeda W, Itoh S, Deguchi-Tawarada M, Ohtsuka T, Mizoguchi A, Takai Y
Journal of cell science (2005) 118(Pt 6): 1267-77.
- SynCAM, a synaptic adhesion molecule that drives synapse assembly.
Biederer T, Sara Y, Mozhayeva M, Atasoy D, Liu X, Kavalali ET, Südhof TC
Science (New York, N.Y.) (2002) 297(5586): 1525-31.

TO BE USED IN VITRO / FOR RESEARCH ONLY

NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS

Development of synapses involves transsynaptic interactions of dedicated synaptic adhesion molecules like neuroligins, neurexins and synCAMs (**Synaptic cell adhesion molecules**). Four SynCAM isoforms have been described so far. All share a common domain structure and contain three extracellular Ig domains, a single transmembrane region, and a short COOH-terminal cytoplasmic tail. **SynCAM 2** has been shown to be associated to myelinated axons and **synCAM 3** can serve as a receptor for herpes viruses.