



Rudolf-Wissell-Str. 28  
37079 Göttingen, Germany  
Phone: +49 551-50556-0  
Fax: +49 551-50556-384  
E-mail: sales@sysy.com  
Web: www.sysy.com

## SynCAM 4

Cat.No. 243 403; Polyclonal rabbit antibody, 50 µg specific antibody (lyophilized)

### Data Sheet

Reconstitution/ Storage	50 µg specific antibody, lyophilized. Affinity purified with the immunogen. Rabbit serum albumin was added for stabilization. For reconstitution add 50 µl H <sub>2</sub> O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C until use.
Applications	<b>WB:</b> 1 : 1000 (AP staining) <b>IP:</b> not tested yet <b>ICC:</b> 1 : 100 up to 1 : 500 <b>IHC:</b> not tested yet <b>IHC-P/FFPE:</b> not tested yet
Immunogen	Synthetic peptide corresponding to AA 376 to 388 from mouse SynCAM4
Reactivity	Reacts with: rat (Q1WIM1), mouse (Q8R464). Other species not tested yet.
Specificity matching control	Specific for synCAM 4. 243-4P

### Selected General References

- Aberrations of a cell adhesion molecule CADM4 in renal clear cell carcinoma. Nagata M, Sakurai-Yageta M, Yamada D, Goto A, Ito A, Fukuhara H, Kume H, Morikawa T, Fukayama M, Homma Y, Murakami Y, et al. International journal of cancer (2012) 130(6): 1329-37.
- Localization of Cadm2a and Cadm3 proteins during development of the zebrafish nervous system. Hunter PR, Nikolaou N, Odermatt B, Williams PR, Drescher U, Meyer MP. The Journal of comparative neurology (2011) 519(11): 2252-70.
- The cell adhesion nectin-like molecules (Necl) 1 and 4 suppress the growth and tumorigenic ability of colon cancer cells. Raveh S, Gavert N, Spiegel I, Ben-Ze'ev A. Journal of cellular biochemistry (2009) 108(1): 326-36.
- The adhesion molecule Necl-3/SynCAM-2 localizes to myelinated axons, binds to oligodendrocytes and promotes cell adhesion. Pellissier F, Gerber A, Bauer C, Ballivet M, Ossipow V. BMC neuroscience (2007) 8: 90.
- Nectin-like proteins mediate axon Schwann cell interactions along the internode and are essential for myelination. Maurel P, Einheber S, Galinska J, Thaker P, Lam I, Rubin MB, Scherer SS, Murakami Y, Gutmann DH, Salzer JL. The Journal of cell biology (2007) 178(5): 861-74.
- SynCAMs organize synapses through heterophilic adhesion. Fogel Al, Akins MR, Krupp AJ, Stagi M, Stein V, Biederer T. The Journal of neuroscience : the official journal of the Society for Neuroscience (2007) 27(46): 12516-30.
- Crystal structure of the V domain of human Nectin-like molecule-1/Syncam3/Tsl1/Igsf4b, a neural tissue-specific immunoglobulin-like cell-cell adhesion molecule. Dong X, Xu F, Gong Y, Gao J, Lin P, Chen T, Peng Y, Qiang B, Yuan J, Peng X, Rao Z, et al. The Journal of biological chemistry (2006) 281(15): 10610-7.
- Nectin-like molecule-1/TSLL1/SynCAM3: a neural tissue-specific immunoglobulin-like cell-cell adhesion molecule localizing at non-junctional contact sites of presynaptic nerve terminals, axons and glia cell processes. Kakunaga S, Ikeda W, Itoh S, Deguchi-Tawarada M, Ohtsuka T, Mizoguchi A, Takai Y. Journal of cell science (2005) 118(Pt 6): 1267-77.
- SynCAM, a synaptic adhesion molecule that drives synapse assembly. Biederer T, Sara Y, Mozhayeva M, Atasoy D, Liu X, Kavalali ET, Südhof TC. Science (New York, N.Y.) (2002) 297(5586): 1525-31.

### TO BE USED IN VITRO / FOR RESEARCH ONLY

NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS

Development of synapses involves transsynaptic interactions of dedicated synaptic adhesion molecules like neuroligins, neurexins and synCAMs (**Synaptic cell adhesion molecules**). Four SynCAM isoforms have been described so far. All share a common domain structure and contain three extracellular Ig domains, a single transmembrane region, and a short COOH-terminal cytoplasmic tail. **SynCAM 2** has been shown to be associated to myelinated axons and **synCAM 3** can serve as a receptor for herpes viruses.