

## CD 3e

Cat.No. 413 103; Polyclonal rabbit antibody, 50 µg specific antibody (lyophilized)

### Data Sheet

Reconstitution/ Storage	50 µg specific antibody, lyophilized. Affinity purified with the immunogen. Rabbit serum albumin was added for stabilization. For reconstitution add 50 µl H <sub>2</sub> O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C until use.
Applications	<b>WB:</b> not tested yet <b>IP:</b> not tested yet <b>ICC:</b> not tested yet <b>IHC:</b> not tested yet <b>IHC-P/FFPE:</b> 1 : 1000
Immunogen	Synthetic peptide corresponding to AA 44 to 64 from mouse CD3e (UniProt Id: P22646)
Reactivity	Reacts with: mouse (P22646). No signal: human (P07766), rat (D4A5M2). Other species not tested yet.
Specificity	Specific for mouse CD 3e without cross-reactivity to human and rat CD 3e.

**TO BE USED IN VITRO / FOR RESEARCH ONLY**  
**NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS**

Cluster of differentiation 3 (CD3) is a defining feature of cells belonging to the T cell lineage. It is composed of the four subunits CD3 gamma, CD3 delta, **CD3 epsilon (CD3e)** and CD3 zeta, that form a multimeric protein complex. This complex associates with the T cell receptor (TCR) and serves as a T cell co-receptor. The CD3 molecules contain immunoreceptor tyrosine-based activation motifs (ITAMs) that serve as the nucleating point for the intracellular signal transduction machinery upon TCR engagement. TCR/CD3 signaling is central to the initiation of antigen-specific T cell responses to pathogens and vaccines, as well as transplanted tissues, tumors, and autoantigens. CD3 is initially expressed in the cytoplasm of pro-thymocytes. During T cell maturation the expression of CD3 migrates to the cell-membrane. The specific appearance at all stages of T cell development make CD3 a useful immunohistochemical marker for T cells in tissue sections. In the clinical setting, CD3 is a relevant marker for the classification of malignant lymphomas and leukemias as the antigen remains present in almost all T-cell lymphomas and leukemias. It can also be used to detect T cells in celiac disease, lymphocytic and collagenous colitis.

### Selected General References

CD3 immunohistochemical staining in diagnosis of lymphocytic colitis.  
Fiehn AM, Engel U, Holck S, Munck LK, Engel PJ  
Human pathology (2016) 48: 25-31.

T cell activation.  
Smith-Garvin JE, Koretzky GA, Jordan MS  
Annual review of immunology (2009) 27: 591-619.

Lymphocytic and collagenous colitis: an immunohistochemical study.  
Mosnier JF, Larvol L, Barge J, Dubois S, De La Bigne G, Hénin D, Cerf M  
The American journal of gastroenterology (1996) 91(4): 709-13.

CD3: structure, function, and role of immunostaining in clinical practice.  
Chetty R, Gatter K  
The Journal of pathology (1994) 173(4): 303-7.