

Rudolf-Wissell-Str. 28 37079 Göttingen, Germany

Phone: +49 551-50556-0
Fax: +49 551-50556-384
E-mail: sales@sysy.com
Web: www.sysy.com

Synaptogyrin 2

Cat.No. 103 203; Polyclonal rabbit antibody, 50 µg specific antibody (lyophilized)

Data Sheet

Reconstitution/ Storage	50 μg specific antibody, lyophilized. Affinity purified with the immunogen. Rabbit serum albumin was added for stabilization. For reconstitution add 50 μl H $_2$ O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C until use.
Applications	WB: 1: 1000 up to 1: 2000 AP-staining IP: not tested yet ICC: not tested yet IHC: not tested yet IHC-P/FFPE: not tested yet
Immunogen	Synthetic peptide corresponding to AA 211 to 224 from rat Synaptogyrin2 (UniProt Id: O54980)
Reactivity	Reacts with: rat (O54980), mouse (O55101). Other species not tested yet.
Specificity	Specific for Synaptogyrin 2.
matching control	103-02P

TO BE USED IN VITRO / FOR RESEARCH ONLY NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS

Synaptogyrins are tyrosine-phosphorylated proteins with two neuronal (synaptogyrin 1 and 3) and one ubiquitous, **synaptogyrin 2** or **cellugyrin** isoform.

Synaptogyrins are integral membrane proteins and localize to the membrane of small vesicles. Synaptogyrin 1 and 3 are expressed in the brain whereby the latter shows a more restricted expression pattern with high levels in the mossy fiber region of the hippocampus, substantia nigra pars reticulata, pallidum, and deep cerebellar nuclei.

Synaptogyrin 2/cellugyrin, a close relative, is expressed in all tissues, for instance, in distinct populations of GLUT 4 containing vesicles.

Selected General References

Cellugyrin induces biogenesis of synaptic-like microvesicles in PC12 cells. Belfort GM, Bakirtzi K, Kandror KV

The Journal of biological chemistry (2005) 280(8): 7262-72.

Cellugyrin is a marker for a distinct population of intracellular Glut4-containing vesicles. Kupriyanova TA, Kandror KV

The Journal of biological chemistry (2000) 275(46): 36263-8.

Cellugyrin, a novel ubiquitous form of synaptogyrin that is phosphorylated by pp60c-src. Janz R. Südhof TC

The Journal of biological chemistry (1998) 273(5): 2851-7.