

Cytohesin 2

Cat.No. 212 203; Polyclonal rabbit antibody, 50 µg specific antibody (lyophilized)

Data Sheet

Reconstitution/ Storage	50 µg specific antibody, lyophilized. Affinity purified with the immunogen. Rabbit serum albumin was added for stabilization. For reconstitution add 50 µl H ₂ O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C until use.
Applications	WB: 1 : 1000 up to 1 : 5000 (AP staining) IP: yes ICC: not tested yet IHC: not tested yet IHC-P/FFPE: not tested yet
Immunogen	Recombinant protein corresponding to AA 1 to 65 from rat Cytohesin2 (UniProt Id: P63035)
Reactivity	Reacts with: rat (P97696), mouse (O08967). Other species not tested yet.
Specificity	Specific for cytohesin 2.

TO BE USED IN VITRO / FOR RESEARCH ONLY
NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS

Cytohesins are a group of guanine nucleotide exchange factors (GEFs) that contain an N-terminal coil, a central Sec-7 and a C-terminal pleckstrin homology (ph) domain.

They control the activity of ADP-ribosylation factors (Arfs) and are involved in the control of vesicular transport and organization of the actin cytoskeleton.

Cytohesins have also been reported to bind to integrins, a diverse family of heterodimeric transmembrane adhesion receptors that are present on most vertebrate cell types. These proteins are known to be involved in cell-cell or cell-matrix interactions.

Selected General References

ARL4D recruits cytohesin-2/ARNO to modulate actin remodeling.

Li CC, Chiang TC, Wu TS, Pacheco-Rodriguez G, Moss J, Lee FJ

Molecular biology of the cell (2007) 18(11): 4420-37.

Actin cytoskeletal association of cytohesin-1 is regulated by specific phosphorylation of its carboxyl-terminal polybasic domain.

Dierks H, Kolanus J, Kolanus W

The Journal of biological chemistry (2001) 276(40): 37472-81.

Similarities in function and gene structure of cytohesin-4 and cytohesin-1, guanine nucleotide-exchange proteins for ADP-ribosylation factors.

Ogasawara M, Kim SC, Adamik R, Togawa A, Ferrans VJ, Takeda K, Kirby M, Moss J, Vaughan M

The Journal of biological chemistry (2000) 275(5): 3221-30.

Solution structure of the cytohesin-1 (B2-1) Sec7 domain and its interaction with the GTPase ADP ribosylation factor 1.

Betz SF, Schnuchel A, Wang H, Olejniczak ET, Meadows RP, Lipsky BP, Harris EA, Staunton DE, Fesik SW

Proceedings of the National Academy of Sciences of the United States of America (1998) 95(14): 7909-14.

Cytohesin-1, a cytosolic guanine nucleotide-exchange protein for ADP-ribosylation factor.

Meacci E, Tsai SC, Adamik R, Moss J, Vaughan M

Proceedings of the National Academy of Sciences of the United States of America (1997) 94(5): 1745-8.