

PSD 95

Cat.No. 124 003; Polyclonal rabbit antibody, 50 µg specific antibody (lyophilized)

Data Sheet

Reconstitution/ Storage	50 µg specific antibody, lyophilized. Affinity purified with the immunogen. Rabbit serum albumin was added for stabilization. For reconstitution add 50 µl H ₂ O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C until use.
Applications	WB: 1 : 1000 up to 1 : 2000 (AP staining) IP: not tested yet ICC: 1 : 500 IHC: not tested yet IHC-P/FFPE: not tested yet
Immunogen	Synthetic peptide corresponding to AA 18 to 32 from rat PSD95 (UniProt Id: P31016)
Reactivity	Reacts with: rat (P31016), mouse (Q62108). Other species not tested yet.
Specificity	Specific for PSD 95. (K.O. verified)
matching control	124-0P

TO BE USED IN VITRO / FOR RESEARCH ONLY

NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS

PSD 95 (postsynaptic density protein 95 kDa, also called **SAP 90**: synapse associated protein of 90 kDa and **DLG 4**) is a component of postsynaptic densities in central synapses. It contains three PDZ domains. The first and second PDZ domain localizes NMDA receptors and K⁺ channels to synapses, the third binds to neuroligins which are neuronal cell adhesion molecules that interact with β-neurexins and form intercellular junctions. Thus different PDZ domains of PSD 95 might be specialized for distinct functions.

Selected References SY SY Antibodies

Kibra Modulates Learning and Memory via Binding to Dendrin.
 Ji Z, Li H, Yang Z, Huang X, Ke X, Ma S, Lin Z, Lu Y, Zhang M
Cell reports (2019) 26(8): 2064-2077.e7. **ICC; tested species: mouse**

Selected General References

- SAP family proteins.
 Fujita A, Kurachi Y
Biochemical and biophysical research communications (2000) 269(1): 1-6.
- Molecular organization of excitatory chemical synapses in the mammalian brain.
 Gundelfinger ED, tom Dieck S
Die Naturwissenschaften (2000) 87(12): 513-23.
- Binding of neuroligins to PSD-95.
 Irie M, Hata Y, Takeuchi M, Ichchenko K, Toyoda A, Hirao K, Takai Y, Rosahl TW, Südhof TC
Science (New York, N.Y.) (1997) 277(5331): 1511-5.
- Mechanisms determining the time course of secretion in neuroendocrine cells.
 Chow RH, Klingauf J, Heinemann C, Zucker RS, Neher E
Neuron (1996) 16(2): 369-76.
- Domain interaction between NMDA receptor subunits and the postsynaptic density protein PSD-95.
 Kornau HC, Schenker LT, Kennedy MB, Seuberg PH
Science (New York, N.Y.) (1995) 269(5231): 1737-40.