

Dynamin 3

Cat.No. 115 303; Polyclonal rabbit antibody, 50 µg specific antibody (lyophilized)

Data Sheet

Reconstitution/ Storage	50 µg specific antibody, lyophilized. Affinity purified with the immunogen. Rabbit serum albumin was added for stabilization. For reconstitution add 50 µl H ₂ O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C until use.
Applications	WB: 1 : 1000 up to 1 : 5000 (AP staining) IP: yes ICC: 1 : 500 IHC: 1 : 500 IHC-P/FFPE: 1 : 500
Immunogen	Synthetic peptide corresponding to AA 773 to 794 from mouse Dynamin3 (UniProt Id: Q8BZ98)
Reactivity	Reacts with: rat (Q08877), mouse (Q8BZ98). Other species not tested yet.
Specificity	Specific for dynamin 3.
matching control	115-3P

Selected General References

Dynamin and its role in membrane fission.
 Hinshaw JE
Annual review of cell and developmental biology (2000) 16: 483-519.

Accessory factors in clathrin-dependent synaptic vesicle endocytosis.
 Slepnev VI, De Camilli P
Nature reviews. Neuroscience (2000) 1(3): 161-72.

Sequential steps in clathrin-mediated synaptic vesicle endocytosis.
 Brodin L, Löw P, Shupliakov O
Current opinion in neurobiology (2000) 10(3): 312-20.

Synaptic vesicle biogenesis.
 Hannah MJ, Schmidt AA, Huttner WB
Annual review of cell and developmental biology (1999) 15: 733-98.

The synaptic vesicle cycle: a cascade of protein-protein interactions.
 Südhof TC
Nature (1995) 375(6533): 645-53.

Complexins: cytosolic proteins that regulate SNAP receptor function.
 McMahon HT, Missler M, Li C, Südhof TC
Cell (1995) 83(1): 111-9.

Synaptic vesicles and exocytosis.
 Jahn R, Südhof TC
Annual review of neuroscience (1994) 17: 219-46.

Differential expression and regulation of multiple dynamins.
 Sontag JM, Fykse EM, Ushkaryov Y, Liu JP, Robinson PJ, Südhof TC
The Journal of biological chemistry (1994) 269(6): 4547-54.

Dynamin GTPase regulated by protein kinase C phosphorylation in nerve terminals.
 Robinson PJ, Sontag JM, Liu JP, Fykse EM, Slaughter C, McMahon H, Südhof TC
Nature (1993) 365(6442): 163-6.

Molecular cloning of the microtubule-associated mechanochemical enzyme dynamin reveals homology with a new family of GTP-binding proteins.
 Obar RA, Collins CA, Hammarback JA, Shpetner HS, Vallee RB
Nature (1990) 347(6290): 256-61.

**TO BE USED IN VITRO / FOR RESEARCH ONLY
 NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS**

Dynamin was discovered because of its binding to microtubules. It was later shown not to function in the cytoskeleton but in endocytosis. Dynamin is required for clathrin - mediated endocytosis. It contains a NH2 - terminal GTPase domain, a middle pleckstrin - homology domain, and a COOH-terminal proline - rich sequence. The COOH - terminal sequence binds to amphiphilin which contains a SH3 domain that recognizes the proline - rich sequence of dynamin. There are at least three isoforms of dynamin: Dynamin 1 is enriched in synapses whereas dynamin 2 is ubiquitous and dynamin 3 is expressed in brain and testis. Neuronal dynamin 1 is phosphorylated by protein kinase C and dephosphorylated by calcineurin during an action potential in the nerve terminal. It is possible that the dephosphorylation provides a trigger for endocytosis.