

GFP

Cat.No. 132 003; Polyclonal rabbit antibody, 50 µg specific antibody (lyophilized)

Data Sheet

Reconstitution/ Storage	50 µg specific antibody, lyophilized. Affinity purified with the immunogen. Rabbit serum albumin was added for stabilization. For reconstitution add 50 µl H ₂ O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C until use.
Applications	WB: 1 : 1000 up to 1 : 20000 (AP staining) IP: yes ICC: 1 : 500 IHC: 1 : 500 IHC-P/FFPE: not tested yet
Immunogen	Recombinant protein corresponding to AA 1 to 238 from GFP (UniProt Id: P42212)
Specificity	Recognizes GFP, mEGFP, superfolder GFP, most common CFP and YFP variants. Does not cross-react to mCherry, mRFP, dsRed, mTagBFP or their most common derivatives.

TO BE USED IN VITRO / FOR RESEARCH ONLY NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS

Green fluorescent protein **GFP** and its derivatives have become universal tools in cell biology. These antibodies allow immunoprecipitation and visualization of GFP fusion proteins on immunoblots and by immunocytochemistry.

Selected References SYSY Antibodies

Tyrosine phosphorylation regulates the endocytosis and surface expression of GluN3A-containing NMDA receptors. Chowdhury D, Marco S, Brooks IM, Zanduetta A, Rao Y, Haucke V, Wesseling JF, Tavalin SJ, Pérez-Otaño I. The Journal of neuroscience : the official journal of the Society for Neuroscience (2013) 33(9): 4151-64. **IP, ICC**

The catalytic function of the gephyrin-binding protein IQSEC3 regulates neurotransmitter-specific matching of pre- and postsynaptic structures in primary hippocampal cultures. Früh S, Tyagarajan SK, Campbell B, Bosshard G, Fritschy JM. Journal of neurochemistry (2018) : . **WB; tested species: rat**

Drosophila gene tao-1 encodes proteins with and without a Ste20 kinase domain that affect cytoskeletal architecture and cell migration differently. Pflanz R, Voigt A, Yakulov T, Jäckle H. Open biology (2015) 5(1): 140161. **IHC**

Ca²⁺/calmodulin binding to PSD-95 mediates homeostatic synaptic scaling down. Chowdhury D, Turner M, Patriarchi T, Hergarden AC, Anderson D, Zhang Y, Sun J, Chen CY, Ames JB, Hell JW. The EMBO journal (2018) 37(1): 122-138. **IP**

Revisiting adult neurogenesis and the role of erythropoietin for neuronal and oligodendroglial differentiation in the hippocampus. Hassouna I, Ott C, Wüstefeld L, Offen N, Neher RA, Mitkovski M, Winkler D, Sperling S, Fries L, Goebbels S, Vreja IC, et al. Molecular psychiatry (2016) 21(12): 1752-1767. **IHC**

The GTPase Rab26 links synaptic vesicles to the autophagy pathway. Binotti B, Pavlos NJ, Riedel D, Wenzel D, Vorbrüggen G, Schalk AM, Kühnel K, Boyken J, Erck C, Martens H, Chua JJ, et al. eLife (2015) 4: e05597. **IP**

Bällchen is required for self-renewal of germline stem cells in Drosophila melanogaster. Herzig B, Yakulov TA, Klinge K, Günesdogan U, Jäckle H, Herzig A. Biology open (2014) 3(6): 510-21. **ICC**

Morphological, biophysical and synaptic properties of glutamatergic neurons of the mouse spinal dorsal horn. Punnakkal P, von Schoultz C, Haenraets K, Wildner H, Zeilhofer HU. The Journal of physiology (2014) 592(4): 759-76. **IHC**

Titration of Syntaxin1 in mammalian synapses reveals multiple roles in vesicle docking, priming, and release probability. Arancillo M, Min SW, Gerber S, Münster-Wandowski A, Wu YJ, Herman M, Trimbuch T, Rah JC, Ahnert-Hilger G, Riedel D, Südhof TC, et al. The Journal of neuroscience : the official journal of the Society for Neuroscience (2013) 33(42): 16698-714. **ICC**

Oligodendrocyte dynamics in the healthy adult CNS: evidence for myelin remodeling. Young KM, Psachoulia K, Tripathi RB, Dunn SJ, Cossell L, Attwell D, Tohyama K, Richardson WD. Neuron (2013) 77(5): 873-85. **IHC; tested species: mouse**

Selective regulation of NR2B by protein phosphatase-1 for the control of the NMDA receptor in neuroprotection. Farinelli M, Heitz FD, Grewe BF, Tyagarajan SK, Helmchen F, Mansuy IM. PLoS one (2012) 7(3): e34047. **IHC**

The short-time structural plasticity of dendritic spines is altered in a model of Rett syndrome. Landi S, Putignano E, Boggio EM, Giustetto M, Pizzorusso T, Ratto GM. Scientific reports (2011) 1: 45. **IHC; tested species: mouse**

Protein kinase C activation promotes the internalization of the human cationic amino acid transporter hCAT-1. A new regulatory mechanism for hCAT-1 activity. Rotmann A, Strand D, Martiné U, Closs EI. The Journal of biological chemistry (2004) 279(52): 54185-92. **IP**

Selected General References

Imaging into the future: visualizing gene expression and protein interactions with fluorescent proteins. van Roesel P, Brand AH. Nature cell biology (2002) 4(1): E15-20.

Illuminating the secretory pathway: when do we need vesicles? Stephens DJ, Pepperkok R. Journal of cell science (2001) 114(Pt 6): 1053-9.

Watching proteins in the wild: fluorescence methods to study protein dynamics in living cells. Chamberlain C, Hahn KM. Traffic (Copenhagen, Denmark) (2000) 1(10): 755-62.