

Cat.No. 170 003; Polyclonal rabbit antibody, 50 µg specific antibody (lyophilized)

Data Sheet

| | |
|-------------------------|---|
| Reconstitution/ Storage | 50 µg specific antibody, lyophilized. Affinity purified with the immunogen. Rabbit serum albumin was added for stabilization. For reconstitution add 50 µl H ₂ O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C until use. |
| Applications | WB: 1 : 1000 (AP staining) IP: yes ICC: 1 : 100 up to 1 : 1000 IHC: not tested yet IHC-P/FFPE: not tested yet |
| Immunogen | Recombinant protein corresponding to AA 2 to 190 from rat Membrin (UniProt Id: O35165) |
| Reactivity | Reacts with: rat (O35165), mouse (O35166). Other species not tested yet. |
| Specificity | Specific for membrin. |

TO BE USED IN VITRO / FOR RESEARCH ONLY NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS

Membrin also known as **GOSR 2, GS 27** and **GOS 27** is a 27 kDa membrane protein on the surface of the Golgi compartment. It has been shown to serve as a tSNARE in ER to Golgi transport. In co-immunoprecipitation studies membrin was identified as a member of a SNARE complex consisting of syntaxin 5, GOSR 1 (GS 28, GOS 28), rbet1 and rsec22. A more detailed analysis revealed two subcomplexes within this complex. One contains syntaxin 5 (mainly the shorter 35 kDa variant) and GS 28 whereas the other is composed of syntaxin 5 (35 and 42 kDa variant), membrin, rsec22 and rbet1. It has been suggested that these complexes mediate the fusion of ER-derived vesicles with vesicular tubular clusters (VTC), and the fusion of VTCs to form the cis-Golgi compartment.

Selected References SYSY Antibodies

COG6 interacts with a subset of the Golgi SNAREs and is important for the Golgi complex integrity. Kudlyk T, Willett R, Pokrovskaya ID, Lupashin V. *Traffic* (Copenhagen, Denmark) (2013) 14(2): 194-204. **WB**

Selected General References

Targeting of Arf-1 to the early Golgi by membrin, an ER-Golgi SNARE. Honda A, Al-Awar OS, Hay JC, Donaldson JG. *The Journal of cell biology* (2005) 168(7): 1039-51.

Countercurrent distribution of two distinct SNARE complexes mediating transport within the Golgi stack. Volchuk A, Ravazzola M, Perrelet A, Eng WS, Di Liberto M, Varlamov O, Fukasawa M, Engel T, Söllner TH, Rothman JE, Orci L, et al. *Molecular biology of the cell* (2004) 15(4): 1506-18.

The SNARE motif contributes to rbet1 intracellular targeting and dynamics independently of SNARE interactions. Joglekar AP, Xu D, Rigotti DJ, Fairman R, Hay JC. *The Journal of biological chemistry* (2003) 278(16): 14121-33.

Sequential tethering of Golgins and catalysis of SNAREpin assembly by the vesicle-tethering protein p115. Shorter J, Beard MB, Seemann J, Dirac-Svejstrup AB, Warren G. *The Journal of cell biology* (2002) 157(1): 45-62.

Subunit structure of a mammalian ER/Golgi SNARE complex. Xu D, Joglekar AP, Williams AL, Hay JC. *The Journal of biological chemistry* (2000) 275(50): 39631-9.

SNARE membrane trafficking dynamics in vivo. Chao DS, Hay JC, Winnick S, Prekeris R, Klumperman J, Scheller RH. *The Journal of cell biology* (1999) 144(5): 869-81.

Localization, dynamics, and protein interactions reveal distinct roles for ER and Golgi SNAREs. Hay JC, Klumperman J, Oorschot V, Steegmaier M, Kuo CS, Scheller RH. *The Journal of cell biology* (1998) 141(7): 1489-502.

Role of vesicle-associated syntaxin 5 in the assembly of pre-Golgi intermediates. Rowe T, Dascher C, Bannykh S, Plutner H, Balch WE. *Science (New York, N.Y.)* (1998) 279(5351): 696-700.