

Histidine Decarboxylase

Cat.No. 405 003; Polyclonal rabbit antibody, 50 µg specific antibody (lyophilized)

Data Sheet

Reconstitution/ Storage	50 µg specific antibody, lyophilized. Affinity purified with the immunogen. Rabbit serum albumin was added for stabilization. For reconstitution add 50 µl H ₂ O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C until use.
Applications	WB: not tested yet IP: not tested yet ICC: not tested yet IHC: 1 : 500 IHC-P/FFPE: 1 : 200
Immunogen	Recombinant protein corresponding to AA 1 to 658 from rat HDC (UniProt Id: P16453)
Reactivity	Reacts with: mouse (P23738), rat (P16453). Other species not tested yet.
Specificity	Specific of histamine decarboxylase.

TO BE USED IN VITRO / FOR RESEARCH ONLY
NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS

Histamine is synthesized by histidine decarboxylase (**HDC**) in many different cell types. Mast cells and basophil granulocytes store histamine in their granules that is released upon external stimulation. In addition histamine acts as a modulatory neurotransmitter in the mammalian brain. HDC expressing neurons occur in the tuberomammillary nucleus (TMN), which is located in the hypothalamus. It is the only site of neuronal histamine synthesis in the adult mammalian brain.

Selected General References

Granule maturation in mast cells: histamine in control.
Hallgren J, Gurish MF
European journal of immunology (2014) 44(1): 33-6.

The histaminergic network in the brain: basic organization and role in disease.
Panula P, Nuutinen S
Nature reviews. Neuroscience (2013) 14(7): 472-87.

Histamine neurons in the tuberomammillary nucleus: a whole center or distinct subpopulations?
Blandina P, Munari L, Provensi G, Passani MB
Frontiers in systems neuroscience (2012) 6: 33.

Changes in motoric, exploratory and emotional behaviours and neuronal acetylcholine content and 5-HT turnover in histidine decarboxylase-KO mice.
Dere E, De Souza-Silva MA, Spieler RE, Lin JS, Ohtsu H, Haas HL, Huston JP
The European journal of neuroscience (2004) 20(4): 1051-8.

Degradation of the 74 kDa form of L-histidine decarboxylase via the ubiquitin-proteasome pathway in a rat basophilic/mast cell line (RBL-2H3).
Tanaka S, Nemoto K, Yamamura E, Ohmura S, Ichikawa A
FEBS letters (1997) 417(2): 203-7.