

## Homer 1b/c

Cat.No. 160-02P; control protein, 100 µg protein (lyophilized)

### Data Sheet

Reconstitution/	100 µg protein, lyophilized. For reconstitution add 100 µl H <sub>2</sub> O to get a 1mg/ml solution in TBS. Then aliquot and store at -20°C until use.
Storage	
Immunogen	Recombinant protein corresponding to AA 152 to 354 from human Homer1b/c (UniProt Id: Q86YM7-1)
Recommended dilution	Optimal concentrations should be determined by the end-user.
matching antibodies	160 022, 160 023
Remarks	This control protein consists of the recombinant protein (aa 152-354 of human homer 1b) that has been used for immunization. It has been tested in preadsorption experiments and blocks efficiently and specifically the corresponding signal in Western blots. The amount of protein needed for efficient blocking depends on the titer and on the affinity of the antibody to the antigen.

**TO BE USED IN VITRO / FOR RESEARCH ONLY  
NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS**

Homer is a scaffolding protein of the post synaptic density (PSD) and enriched at excitatory synapses. The protein binds metabotropic glutamate receptors, TRPC1, proteins of the Shank family and others. By aggregating these proteins into clusters, homer was suggested to organize distinct signalling domains.

Three isoforms, **Homer 1**, 2 and 3 have been described. Each of these isoforms is subject to alternative splicing yielding the splice variants a, b, c, d.

### Selected General References

Surface clustering of metabotropic glutamate receptor 1 induced by long Homer proteins.  
Kammermeier PJ  
BMC neuroscience (2006) 7: 1.

Homer 1a enhances spike-induced calcium influx via L-type calcium channels in neocortex pyramidal cells.  
Yamamoto K, Sakagami Y, Sugiura S, Inokuchi K, Shimohama S, Kato N  
The European journal of neuroscience (2005) 22(6): 1338-48.

Differential expression of Homer family proteins in the developing mouse brain.  
Shiraishi Y, Mizutani A, Yuasa S, Mikoshiba K, Furuchi T  
The Journal of comparative neurology (2004) 473(4): 582-99.

Modulation of synaptic signalling complexes by Homer proteins.  
Thomas U  
Journal of neurochemistry (2002) 81(3): 407-13.

Homer-dependent cell surface expression of metabotropic glutamate receptor type 5 in neurons.  
Ango F, Robbe D, Tu JC, Xiao B, Worley PF, Pin JP, Bockaert J, Fagni L  
Molecular and cellular neurosciences (2002) 20(2): 323-9.

An N-terminal sequence specific for a novel Homer1 isoform controls trafficking of group I metabotropic glutamate receptor in mammalian cells.  
Saito H, Kimura M, Inanobe A, Ohe T, Kurachi Y  
Biochemical and biophysical research communications (2002) 296(3): 523-9.

Regulation of dendritic spine morphology and synaptic function by Shank and Homer.  
Sala C, Piéch V, Wilson NR, Passafaro M, Liu G, Sheng M  
Neuron (2001) 31(1): 115-30.

Homer-1c/Vesl-1L modulates the cell surface targeting of metabotropic glutamate receptor type 1alpha: evidence for an anchoring function.  
Ciruela F, Soloviev MM, Chan WY, McIlhinney RA  
Molecular and cellular neurosciences (2000) 15(1): 36-50.

Homer: a link between neural activity and glutamate receptor function.  
Xiao B, Tu JC, Worley PF  
Current opinion in neurobiology (2000) 10(3): 370-4.

Molecular characterisation of two structurally distinct groups of human homers, generated by extensive alternative splicing.  
Soloviev MM, Ciruela F, Chan WY, McIlhinney RA  
Journal of molecular biology (2000) 295(5): 1185-200.

Coupling of mGluR/Homer and PSD-95 complexes by the Shank family of postsynaptic density proteins.  
Tu JC, Xiao B, Naïsbit S, Yuan JP, Petralia RS, Brakeman P, Doan A, Aakalu VK, Lanahan AA, Sheng M, Worley PF, et al.  
Neuron (1999) 23(3): 583-92.

Homer: protein that selectively binds metabotropic glutamate receptors.  
Brakeman PR, Lanahan AA, O'Brien R, Roche K, Barnes CA, Huganir RL, Worley PF  
Nature (1997) 386(6622): 284-8.