

RFP

Rudolf-Wissell-Str. 28 37079 Göttingen, Germany

Phone: +49 551-50556-0
Fax: +49 551-50556-384
E-mail: sales@sysy.com
Web: www.sysy.com

Cat.No. 390 004; Polyclonal Guinea pig antibody, 100 µl antiserum (lyophilized)

Data Sheet

Reconstitution/ Storage	100 μl antiserum, lyophilized. For reconstitution add 100 μl H_2O , then aliquot and store at -20°C until use.
Applications	WB: 1: 1000 (AP staining) IP: not tested yet ICC: 1: 500 up to 1: 1000 IHC: 1: 200 up to 1: 500 IHC-P/FFPE: 1: 200
Immunogen	Recombinant protein corresponding to AA 1 to 236 from mCherry (UniProt Id: X5DSL3)
Epitop	Epitop: AA 1 to 236 from mCherry (UniProt Id: X5DSL3)
Specificity	Recognizes mRFP, mCherry, mOrgange2, dsRed, tdTomato, mScarlet.

TO BE USED IN VITRO / FOR RESEARCH ONLY NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS

Red fluorescent protein RFP and its derivates have become universal tools in cell biology. Most RFPs derive from a protein isolated from Discosoma sp. They are used as fluorescent tags to investigate expression levels, patterns and protein localization.

Selected General References

Ubiquitous expression of the monomeric red fluorescent protein mCherry in transgenic mice. Fink D, Wohrer S, Pfeffer M, Tombe T, Ong CJ, Sorensen PH Genesis (New York, N.Y.: 2000) (2010) 48(12): 723-9.

Improved monomeric red, orange and yellow fluorescent proteins derived from Discosoma sp. red fluorescent protein. Shaner NC, Campbell RE, Steinbach PA, Giepmans BN, Palmer AE, Tsien RY Nature biotechnology (2004) 22(12): 1567-72.

Diversity and evolution of the green fluorescent protein family.

Labas YA, Gurskaya NG, Yanushevich YG, Fradkov AF, Lukyanov KA, Lukyanov SA, Matz MV

Proceedings of the National Academy of Sciences of the United States of America (2002) 99(7): 4256-61.

Novel fluorescent protein from Discosoma coral and its mutants possesses a unique far-red fluorescence. Fradkov AF, Chen Y, Ding L, Barsova EV, Matz MV, Lukyanov SA FEBS letters (2000) 479(3): 127-30.