

## Neurofilament M

Cat.No. 171 211; Monoclonal mouse antibody, 100 µg purified IgG (lyophilized)

### Data Sheet

Reconstitution/ Storage	100 µg purified IgG, lyophilized. Azide was added before lyophilization. For reconstitution add 100 µl H <sub>2</sub> O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C until use.
Applications	<b>WB:</b> not tested yet <b>IP:</b> not tested yet <b>ICC:</b> 1 : 500 <b>IHC:</b> 1 : 500 up to 1 : 1000 <b>IHC-P/FFPE:</b> 1 : 1000 (see remarks)
Clone	309F11
Subtype	IgG2b (κ light chain)
Immunogen	Recombinant protein corresponding to AA 763 to 848 from mouse Neurofilament M (UniProt Id: P08553)
Epitop	Epitop: AA 763 to 848 from mouse Neurofilament M (UniProt Id: P08553)
Reactivity	Reacts with: mouse (P08553). No signal: rat (P12839). Other species not tested yet.
Specificity	specific for Neurofilament M
Remarks	<b>IHC-P:</b> Cat. no. 171 221 is more sensitive and recommended for this application.

**TO BE USED IN VITRO / FOR RESEARCH ONLY**  
**NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS**

**Neurofilaments** are exclusively expressed in nerve cells and are the major structural component of large-diameter myelinated axons. They are predominately composed of three proteins, Neurofilament H, L and M and are among the most highly phosphorylated neuronal proteins.

### Selected General References

- New movements in neurofilament transport, turnover and disease.  
Barry DM, Millecamps S, Julien JP, Garcia ML  
Experimental cell research (2007) 313(10): 2110-20.
- Regulation between O-GlcNAcylation and phosphorylation of neurofilament-M and their dysregulation in Alzheimer disease.  
Deng Y, Li B, Liu F, Iqbal K, Grundke-Iqbal I, Brandt R, Gong CX  
FASEB journal : official publication of the Federation of American Societies for Experimental Biology (2008) 22(1): 138-45.
- CSF neurofilament proteins in the differential diagnosis of dementia.  
de Jong D, Jansen RW, Pijnenburg YA, van Geel WJ, Borm GF, Kremer HP, Verbeek MM  
Journal of neurology, neurosurgery, and psychiatry (2007) 78(9): 936-8.
- 14-3-3 protein binds to the low molecular weight neurofilament (NFL) mRNA 3' UTR.  
Ge WW, Volkening K, Leystra-Lantz C, Jaffe H, Strong MJ  
Molecular and cellular neurosciences (2007) 34(1): 80-7.
- Differential subcellular localization of phosphorylated neurofilament and tau proteins in degenerating neurons of the human entorhinal cortex.  
Porchet R, Probst A, Dráberová E, Dráber P, Riederer IM, Riederer BM  
Neuroreport (2003) 14(7): 929-33.
- Influence of the axotomy to cell body distance in rat rubrospinal and spinal motoneurons: differential regulation of GAP-43, tubulins, and neurofilament-M.  
Fernandes KJ, Fan DP, Tsui BJ, Cassar SL, Tetzlaff W  
The Journal of comparative neurology (1999) 414(4): 495-510.
- Neurofilament protein is differentially distributed in subpopulations of corticocortical projection neurons in the macaque monkey visual pathways.  
Hof PR, Ungerleider LG, Webster MJ, Gattass R, Adams MM, Sailstad CA, Morrison JH  
The Journal of comparative neurology (1996) 376(1): 112-27.
- Differential dynamics of neurofilament-H protein and neurofilament-L protein in neurons.  
Takeda S, Okabe S, Funakoshi T, Hirokawa N  
The Journal of cell biology (1994) 127(1): 173-85.
- Neurofilament immunoreactivity in myenteric neurons differs from that found in the central nervous system.  
Eaker EY, Shaw G, Sninsky CA  
Gastroenterology (1990) 99(5): 1364-71.
- Intermediate filaments in nervous tissues.  
Liem RK, Yen SH, Salomon GD, Shelanski ML  
The Journal of cell biology (1978) 79(3): 637-45.