



## Neuroligin3 KO lysate

Cat.No. 510-Nlg3-KO; , 100 µg tissue lysate

### Data Sheet

Reconstitution/	100 µg lysate in 1 X SDS PAGE loading buffer 'ready to use'.
Storage	Total volume: 50 µl. Concentration: 2 mg/ml. Aliquot and store at -20°C until use.
Applications	WB: yes (see remarks)
Loading/lane	10-30 µg
Source	Balb-C
Remarks	WB: Since several synaptic proteins aggregate after boiling, these lysates are unboiled. Boil sample before loading, if this is recommended for your protocol.

**TO BE USED IN VITRO / FOR RESEARCH ONLY  
NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS**

Rudolf-Wissell-Str. 28  
37079 Göttingen, Germany  
Phone: +49 551-50556-0  
Fax: +49 551-50556-384  
E-mail: sales@sysy.com  
Web: www.sysy.com

### Selected General References

- Activity-dependent validation of excitatory versus inhibitory synapses by neuroligin-1 versus neuroligin-2.  
Chubykin AA, Atasoy D, Etherton MR, Brose N, Kavalali ET, Gibson JR, Südhof TC  
*Neuron* (2007) 54(6): 919-31.
- Dissection of synapse induction by neuroligins: effect of a neuroligin mutation associated with autism.  
Chubykin AA, Liu X, Comoletti D, Tsigelny I, Taylor P, Südhof TC  
*The Journal of biological chemistry* (2005) 280(23): 22365-74.
- Neuroligin 2 is exclusively localized to inhibitory synapses.  
Varoqueaux F, Jamain S, Brose N  
*European journal of cell biology* (2004) 83(9): 449-56.
- Synaptic targeting of neuroligin is independent of neurexin and SAP90/PSD95 binding.  
Dresbach T, Neeb A, Meyer G, Gundelfinger ED, Brose N  
*Molecular and cellular neurosciences* (2004) 27(3): 227-35.
- Neuroligin 1 is a postsynaptic cell-adhesion molecule of excitatory synapses.  
Song JY, Ichtchenko K, Südhof TC, Brose N  
*Proceedings of the National Academy of Sciences of the United States of America* (1999) 96(3): 1100-5.
- The making of neurexins.  
Missler M, Fernandez-Chacon R, Südhof TC  
*Journal of neurochemistry* (1998) 71(4): 1339-47.
- Structures, alternative splicing, and neurexin binding of multiple neuroligins.  
Ichtchenko K, Nguyen T, Südhof TC  
*The Journal of biological chemistry* (1996) 271(5): 2676-82.
- Neuroligin 1: a splice site-specific ligand for beta-neurexins.  
Ichtchenko K, Hata Y, Nguyen T, Ullrich B, Missler M, Moomaw C, Südhof TC  
*Cell* (1995) 81(3): 435-43.
- The synaptic vesicle cycle: a cascade of protein-protein interactions.  
Südhof TC  
*Nature* (1995) 375(6533): 645-53.