

Rab 3a

Cat.No. 107-1P; control peptide, 100 µg peptide (lyophilized)

Data Sheet

Reconstitution/ Storage	100 µg peptide, lyophilized. For reconstitution add 100 µl H ₂ O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C until use. Control peptides should also be stored at -20°C when still lyophilized!
Immunogen	Synthetic peptide corresponding to AA 2 to 14 from rat Rab3a (UniProt Id: P63012)
Recommended dilution	Optimal concentrations should be determined by the end-user.
matching antibodies	107 102
Remarks	This control peptide consists of the synthetic peptide (ASATDSRYGQKES) that has been used for immunization. It has been tested in preadsorption experiments and blocks efficiently and specifically the corresponding signal in Western blots. The amount of peptide needed for efficient blocking depends on the titer and on the affinity of the antibody to the antigen.

TO BE USED IN VITRO / FOR RESEARCH ONLY NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS

Rab 3 is a member of the Rab protein family that belongs to the ras-related superfamily of small monomeric GTPases. Four related isoforms of Rab 3 are known (**Rab 3a**, **3b**, **3c**, and **3d**). Rab 3a and 3c are predominantly expressed in neurons and neuroendocrine cells where they are localized to synaptic vesicles. Unlike the integral membrane proteins of synaptic vesicles, Rab 3a/c is absent from the Golgi complex and thus does not result in immunostaining of the axo-dendritic region as sometimes seen with e.g. synaptophysin, synaptobrevin/VAMP, or synaptogyrin. Rab 3b and 3d are expressed in non-neuronal tissues such as adipocytes and the exocrine pancreas (3d). It has been shown that overexpression of Rab 3 inhibits Ca²⁺ regulated exocytosis and converts it into an constitutive Ca²⁺ independent exocytosis mechanism.

Selected General References

- RAB3 and synaptotagmin: the yin and yang of synaptic membrane fusion.
Geppert M, Südhof TC
Annual review of neuroscience (1998) 21: 75-95.
- Dominant negative Rab3D mutants reduce GTP-bound endogenous Rab3D in pancreatic acini.
Chen X, Ernst SA, Williams JA
The Journal of biological chemistry (2003) 278(50): 50053-60.
- Rab3D: a regulator of exocytosis in non-neuronal cells.
Millar AL, Pavlos NJ, Xu J, Zheng MH
Histology and histopathology (2002) 17(3): 929-36.
- Molecular cloning of the mouse homologue of Rab3c.
Pavlos NJ, Xu J, Papadimitriou JM, Zheng MH
Journal of molecular endocrinology (2001) 27(1): 117-22.
- The small GTP-binding protein Rab3A regulates a late step in synaptic vesicle fusion.
Geppert M, Goda Y, Stevens CF, Südhof TC
Nature (1997) 387(6635): 810-4.
- Characterization of Rab3A, Rab3B and Rab3C: different biochemical properties and intracellular localization in bovine chromaffin cells.
Lin CG, Lin YC, Liu HW, Kao LS
The Biochemical journal (1997) 324 (Pt 1): 85-90.
- The synaptic vesicle cycle: a cascade of protein-protein interactions.
Südhof TC
Nature (1995) 375(6533): 645-53.
- Synaptic vesicles and exocytosis.
Jahn R, Südhof TC
Annual review of neuroscience (1994) 17: 219-46.
- Localization of Rab5 to synaptic vesicles identifies endosomal intermediate in synaptic vesicle recycling pathway.
Fischer von Mollard G, Stahl B, Walch-Solimena C, Takei K, Daniels L, Khokhlatchev A, De Camilli P, Südhof TC, Jahn R
European journal of cell biology (1994) 65(2): 319-26.
- Rab3C is a synaptic vesicle protein that dissociates from synaptic vesicles after stimulation of exocytosis.
Fischer von Mollard G, Stahl B, Khokhlatchev A, Südhof TC, Jahn R
The Journal of biological chemistry (1994) 269(15): 10971-4.