

Rab 27 B

Cat.No. 168-1P; control protein, 100 µg protein (lyophilized)

Data Sheet

Reconstitution/Storage	100 µg protein, lyophilized. For reconstitution add 100 µl H ₂ O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C until use.
Immunogen	Recombinant protein corresponding to AA 1 to 217 from rat Rab27B (UniProt Id: Q99P74)
Recommended dilution	Optimal concentrations should be determined by the end-user.
matching antibodies	168 103
Remarks	This control protein consists of the recombinant protein that has been used for immunization. It has been tested in preadsorption experiments and blocks efficiently and specifically the corresponding signal in Western blots. The amount of protein needed for efficient blocking depends on the titer and on the affinity of the antibody to the antigen.

TO BE USED IN VITRO / FOR RESEARCH ONLY NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS

Rab 27 proteins are members of the Rab protein family that belongs to the ras-related superfamily of small monomeric GTPases. These proteins are involved in intracellular fusion reactions of vesicles or organelles with their target membranes. Two Rab 27 isoforms, Rab **27A** and **27B**, have been described so far.

Mutations in the Rab 27A gene have been shown to be responsible for the Griscelli syndrome characterized by pigment dilution of the hair and an uncontrolled T-lymphocyte and macrophage activation. This disorder is probably due to the dysfunction of melanosomes in melanocytes and lytic granules in CTLs. Additionally Rab 27A is located on mature insulin granules of pancreatic β-cells and is expressed in the pigment epithelium and choriocapillaris of the retina.

In patients who suffer from Griscelli syndrome because of missense mutations in the Rab 27A gene, Rab 27B is upregulated and partially compensates for Rab 27A dysfunction. Rab 27B also regulates amylase secretion in parotid acinar cells.

Recently it has been shown that Rab 27 is also involved in synaptic transmission in *C. elegans*.

Selected General References

- Regulation of synaptic transmission by RAB-3 and RAB-27 in *Caenorhabditis elegans*. Mahoney TR, Liu Q, Itoh T, Luo S, Hadwiger G, Vincent R, Wang ZW, Fukuda M, Nonet ML. *Molecular biology of the cell* (2006) 17(6): 2617-25.
- Rab7 and Rab27a control two motor protein activities involved in melanosomal transport. Jordens I, Westbroek W, Marsman M, Rocha N, Mommaas M, Huizing M, Lambert J, Naeyaert JM, Neeffjes J. *Pigment cell research* (2006) 19(5): 412-23.
- Rab3A and Rab27A cooperatively regulate the docking step of dense-core vesicle exocytosis in PC12 cells. Tsuboi T, Fukuda M. *Journal of cell science* (2006) 119(Pt 11): 2196-203.
- Functional analysis of Rab27a effector granuphilin in insulin exocytosis. Izumi T, Gomi H, Torii S. *Methods in enzymology* (2005) 403: 216-29.
- Rab27a mediates the tight docking of insulin granules onto the plasma membrane during glucose stimulation. Kasai K, Ohara-Imaizumi M, Takahashi N, Mizutani S, Zhao S, Kikuta T, Kasai H, Nagamatsu S, Gomi H, Izumi T. *The Journal of clinical investigation* (2005) 115(2): 388-96.
- Rab27a: a new face in beta cell metabolism-secretion coupling. Aizawa T, Komatsu M. *The Journal of clinical investigation* (2005) 115(2): 227-30.
- A general role for Rab27a in secretory cells. Tolmachova T, Anders R, Stinchcombe J, Bossi G, Griffiths GM, Huxley C, Seabra MC. *Molecular biology of the cell* (2004) 15(1): 332-44.
- The role of Rab27a in the regulation of melanosome distribution within retinal pigment epithelial cells. Futter CE, Ramalho JS, Jaissle GB, Seeliger MW, Seabra MC. *Molecular biology of the cell* (2004) 15(5): 2264-75.
- Rab27A-binding protein Slp2-a is required for peripheral melanosome distribution and elongated cell shape in melanocytes. Kuroda TS, Fukuda M. *Nature cell biology* (2004) 6(12): 1195-203.
- The small GTPase Rab27B regulates amylase release from rat parotid acinar cells. Imai A, Yoshie S, Nashida T, Shimomura H, Fukuda M. *Journal of cell science* (2004) 117(Pt 10): 1945-53.
- Rab27b is up-regulated in human Griscelli syndrome type II melanocytes and linked to the actin cytoskeleton via exon F-Myosin Va transcripts. Westbroek W, Lambert J, De Schepper S, Kleta R, Van Den Bossche K, Seabra MC, Huizing M, Mommaas M, Naeyaert JM. *Pigment cell research* (2004) 17(5): 498-505.
- Rab27b localizes to zymogen granules and regulates pancreatic acinar exocytosis. Chen X, Li C, Izumi T, Ernst SA, Andrews PC, Williams JA. *Biochemical and biophysical research communications* (2004) 323(4): 1157-62.
- A role for Rab27b in NF-E2-dependent pathways of platelet formation. Tiwari S, Italiano JE, Barral DC, Mules EH, Novak EK, Swank RT, Seabra MC, Shivdasani RA. *Blood* (2003) 102(12): 3970-9.
- Multiple factors contribute to inefficient prenylation of Rab27a in Rab prenylation diseases. Larijani B, Hume AN, Tarafder AK, Seabra MC. *The Journal of biological chemistry* (2003) 278(47): 46798-804.
- Characterization of the molecular defects in Rab27a, caused by RAB27A missense mutations found in patients with Griscelli syndrome. Bahadoran P, Busca R, Chiaverini C, Westbroek W, Lambert J, Bille K, Valony G, Fukuda M, Naeyaert JM, Ortonne JP, Ballotti R, et al. *The Journal of biological chemistry* (2003) 278(13): 11386-92.
- Biochemical and functional characterization of Rab27a mutations occurring in Griscelli syndrome patients. Menasche G, Feldmann J, Houdusse A, Desaymard C, Fischer A, Goud B, de Saint Basile G. *Blood* (2003) 101(7): 2736-42.
- Involvement of Rab27b in the regulated secretion of pituitary hormones. Zhao S, Torii S, Yokota-Hashimoto H, Takeuchi T, Izumi T. *Endocrinology* (2002) 143(5): 1817-24.
- Melanophilin directly links Rab27a and myosin Va through its distinct coiled-coil regions. Nagashima K, Torii S, Yi Z, Igarashi M, Okamoto K, Takeuchi T, Izumi T. *FEBS letters* (2002) 517(1-3): 233-8.