

## RIM 1/2

Cat.No. 140 205; Polyclonal Guinea pig antibody, 50 µg specific antibody (lyophilized)

### Data Sheet

Reconstitution/ Storage	50 µg specific antibody, lyophilized. Affinity purified with the immunogen. Guinea pig serum albumin was added for stabilization. For reconstitution add 50 µl H <sub>2</sub> O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C until use.
Applications	<b>WB:</b> 1 : 1000 (AP staining) <b>IP:</b> not tested yet <b>ICC:</b> 1 : 500 <b>IHC:</b> not tested yet <b>IHC-P/FFPE:</b> not tested yet
Immunogen	Recombinant protein corresponding to AA 1 to 466 from rat Rim2 (UniProt Id: Q9JIS1)
Reactivity	Reacts with: rat (Q9JIS1), mouse (Q9EQZ7). Other species not tested yet.
Specificity	RIM 2 including splice variants, cross reacts to RIM 1.

**TO BE USED IN VITRO / FOR RESEARCH ONLY**  
**NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS**

**RIMs** are presynaptic active zone proteins that regulate Ca<sup>2+</sup> triggered release of neurotransmitters. RIM 1α and RIM 2α are composed of an N-terminal zinc-finger domain, a central PDZ domain and two C-terminal C2 domains that are separated by long alternatively spliced sequences. RIM 2β consists of a specific N-terminus, the central PDZ domain and the C-terminal C2 domains. The mRNA for RIM 2β is transcribed from an internal promoter of the RIM 2α gene. Shorter variants of RIM 2 which comprise only the C-terminal C<sub>2</sub>B domain and some flanking regions are referred to as NIM 2 / RIM 2γ and NIM 3 / RIM 3γ.

### Selected General References

Genomic definition of RIM proteins: evolutionary amplification of a family of synaptic regulatory proteins.  
Wang Y, Südhof TC  
Genomics (2003) 81(2): 126-37.

The RIM/NIM family of neuronal C2 domain proteins. Interactions with Rab3 and a new class of Src homology 3 domain proteins.  
Wang Y, Sugita S, Südhof TC  
The Journal of biological chemistry (2000) 275(26): 20033-44.