
Data Sheet

TagFP Selector

Catalog No.	100 μ l resin/ 200 μ l slurry N0510-S	1000 μ l resin/ 2000 μ l slurry N0510-L
-------------	--	--

Description

TagFP Selector is based on a high-affinity single-domain antibody (sdAb) that is covalently immobilized on 4 % cross-linked agarose beads. The innovative, oriented and selective attachment via a flexible linker guarantees a high accessibility of the sdAbs and largely eliminates batch-to-batch variations. Due to the single-chain nature of sdAbs and their stable and covalent attachment, no leakage of light and heavy chains is observed during elution with SDS sample buffer. TagFP Selector thus features high affinity and superior capacity for corresponding fusion proteins while showing negligible unspecific background.

TagFP Selector is compatible not only with physiological buffers but also with high stringency buffers (see "Buffer Compatibility" below). TagFP Selector thus provides great freedom to adjust the binding and washing conditions to the experimental needs.

For recommended protocols please see our webpage at www.nano-tag.com/protocols.

**To be used *in vitro* / for research only,
not for diagnostic or therapeutic use!**

Non-toxic, non-hazardous, non-infectious.

Properties

Support	4 % cross-linked agarose, bead size 50-150 µm
Coating	sdAb anti-TagFP clone 1H7
Reactivity	<p>Recognizes mTagRFP, mTagBFP, mTagRFP657 and mKate. Reactivity against other TagRFP derivatives not analyzed.</p> <p>Does not cross-react with common GFP- or dsRed/RFP derivatives.</p> <p>For a detailed overview visit our webpage at www.nano-tag.com/protocols.</p>
Capacity	> 4 µg mTagBFP per µl of packed beads
Formulation	50 % slurry in PBS containing 20 % Ethanol
Shipment	Shipped at ambient temperature
Storage	Store at 4 °C, do not freeze
Stability	Stable for 12 months
Buffer	<ul style="list-style-type: none">• Common buffer substances at pH 5 to 9
Compatibility	<ul style="list-style-type: none">• 2% Triton X-100, 1% Tween-20, 1% NP-40, 1% CHAPS, 1% Deoxycholate, 0.1% SDS• 4 M NaCl, 2 M KCl, 1 M MgCl₂, 100 mM EDTA• 4 M urea• 10 mM DTT, 10 mM 2-Mercaptoethanol• RNase A, DNase I, Benzonase, protease inhibitors

For more information please visit our web page at www.nano-tag.com

Limited Use Label License

The purchaser of this product is granted a limited, non-transferable right to use the purchased material only for internal, academic research purposes. The license explicitly excludes the right to resell this product or any of its components. For information on further licensing possibilities please contact info@nano-tag.com

anti-Fluorescent Protein Selector Resins Specificity Chart

Fluorescent Protein	GFP Selector	RFP Selector	TagFP Selector
Sirius	++	-	-
tSapphire	++	-	-
Cerulean	++	-	-
eCFP	++	-	-
mTurquoise2	-	-	-
mTFP (mTeal)	+	-	-
acGFP	+	-	-
EGFP	++	-	-
Emerald GFP	++	-	-
mEGFP (A206K)	++	-	-
mEGFP (L221K)	++	-	-
superecliptic pHluorin	+	-	-
paGFP	+	-	-
superfolder GFP	++	-	-
eYFP	+	-	-
mVenus	++	-	-
Citrine	+	-	-
mOrange2	-	+	-
dsRed1	-	++	-
dsRed2	-	++	-
tdTomato	-	++	-
mRFP	-	+	-
mCherry	-	++	-
mTagBFP	-	-	++
mKate	-	-	++
mTagRFP	-	-	++
mTagRFP657	-	-	++
Dendra2	-	-	-
Dronpa	-	-	-
tdEOS	-	-	-
mEOS3.2	-	-	-

++: strong positive signal; + positive signal; - no signal