

## Synapsin 1/2

Cat.No. 106 003; Polyclonal rabbit antibody, 50 µg specific antibody (lyophilized)

### Data Sheet

Reconstitution/ Storage	50 µg specific antibody, lyophilized. Affinity purified with the immunogen. Rabbit serum albumin was added for stabilization. For reconstitution add 50 µl H <sub>2</sub> O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C until use.
Applications	<b>WB:</b> 1 : 1000 (AP staining) <b>IP:</b> yes <b>ICC:</b> 1 : 500 <b>IHC:</b> 1 : 500 <b>IHC-P/FFPE:</b> 1 : 500 <b>ELISA:</b> yes (see remarks)
Immunogen	Synthetic peptide corresponding to AA 2 to 28 from rat Synapsin1 (UniProt Id: P09951)
Reactivity	Reacts with: human (P17600, Q92777), rat (P09951, Q63537), mouse (O88935, Q64332), hamster, cow, zebrafish. Other species not tested yet.
Specificity	Specific for synapsins 1a/b and 2a/b. (K.O. verified)
matching control	106-0P
Remarks	<b>ELISA:</b> Suitable as detector antibody for sandwich-ELISA with cat. no. 106 001 as capture antibody (protocol for sandwich-ELISA).

### TO BE USED IN VITRO / FOR RESEARCH ONLY NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS

**Synapsins** are neuron-specific phosphoproteins that are exclusively associated with small synaptic vesicles, with little or no expression in other tissues including neuroendocrine cells. In mammals, three distinct synapsin genes (synapsin 1, 2 and 3) encode more than eight neuronal isoforms.

Synapsin 1 is one of the most specific markers of synapses throughout the central and peripheral nervous system. In addition to synaptic nerve terminals, the protein is also present in certain sensory nerve endings. It is expressed in two splice variants (synapsin 1a and synapsin 1b). Synapsin 1 interacts with vesicle membranes as well as with actin and spectrin.

**Synapsin 2** is expressed in the nervous system and also two splice variants were described so far, while synapsin 3 shows a more restricted expression pattern and is mainly found in the hippocampus. Synapsins are major phosphoproteins and are substrates for several protein kinases such as PKA, CaMK I and CaMK II. Synapsin 1 is widely used as reference substrate for calmodulin-dependent protein kinases.

### Selected References SYSY Antibodies

Loss of X-linked Protocadherin-19 differentially affects the behavior of heterozygous female and hemizygous male mice. Hayashi S, Inoue Y, Hattori S, Kaneko M, Shioi G, Miyakawa T, Takeichi M  
Scientific reports (2017) 7(1): 5801. **ICC; tested species: mouse**

Selective Localization of Shanks to VGLUT1-Positive Excitatory Synapses in the Mouse Hippocampus. Heise C, Schroeder JC, Schoen M, Halbedl S, Reim D, Woelfle S, Kreutz MR, Schmeisser MJ, Boeckers TM  
Frontiers in cellular neuroscience (2016) 10: 106. **IHC**

### Selected General References

A phospho-switch controls the dynamic association of synapsins with synaptic vesicles. Hosaka M, Hammer RE, Südhof TC  
Neuron (1999) 24(2): 377-87.

Synapsin-dependent reserve pool of synaptic vesicles supports replenishment of the readily releasable pool under intense synaptic transmission. Vasileva M, Horstmann H, Geumann C, Gitler D, Kuner T  
The European journal of neuroscience (2012) 36(8): 3005-20.

Essential functions of synapsins I and II in synaptic vesicle regulation. Rosahl TW, Spillane D, Missler M, Herz J, Selig DK, Wolff JR, Hammer RE, Malenka RC, Südhof TC  
Nature (1995) 375(6531): 488-93.

The synaptic vesicle cycle: a cascade of protein-protein interactions. Südhof TC  
Nature (1995) 375(6533): 645-53.

Synaptic vesicles and exocytosis. Jahn R, Südhof TC  
Annual review of neuroscience (1994) 17: 219-46.