

Synaptojanin 1

Cat.No. 145-1P; control peptide, 100 µg peptide (lyophilized)

Data Sheet

Reconstitution/ Storage	100 µg peptide, lyophilized. For reconstitution add 100 µl H ₂ O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C until use. Control peptides should also be stored at -20°C when still lyophilized!
Immunogen	Synthetic peptide corresponding to AA 1140 to 1155 from rat Synaptojanin1 (UniProt Id: Q62910-1)
Recommended dilution	Optimal concentrations should be determined by the end-user.
matching antibodies	145 103
Remarks	This control peptide consists of the synthetic peptide (CGVGAPPSPGVTRREME) that has been used for immunization. It has been tested in preadsorption experiments and blocks efficiently and specifically the corresponding signal in Western blots. The amount of peptide needed for efficient blocking depends on the titer and on the affinity of the antibody to the antigen.

TO BE USED IN VITRO / FOR RESEARCH ONLY
NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS

Synaptojanin 1 is a phosphoinositide phosphatase which exists in two tissue specific isoforms (170 and 145 kDa). The 145 kDa isoform is predominantly expressed in the nervous system. It has a three domain structure with an N-terminal part homologous to the yeast Sac 1p protein, a central inositol 5-phosphatase domain and a C-terminal SH3 binding domain. Synaptojanin is involved in clathrin mediated synaptic vesicle recycling and binds to endophilin and amphiphysin. Six isoforms are described so far and two splice variants of the 145 kDa isoform. The predominant one contains a 16 AA insert at position 1140-1155.

Selected General References

- Regulation of synaptojanin 1 by cyclin-dependent kinase 5 at synapses.
Lee SY, Wenk MR, Kim Y, Nairn AC, De Camilli P
Proceedings of the National Academy of Sciences of the United States of America (2004) 101(2): 546-51.
- Endophilin is required for synaptic vesicle endocytosis by localizing synaptojanin.
Schuske KR, Richmond JE, Matthies DS, Davis WS, Runz S, Rube DA, van der Bliek AM, Jorgensen EM
Neuron (2003) 40(4): 749-62.
- Synaptojanin is recruited by endophilin to promote synaptic vesicle uncoating.
Verstreken P, Koh TW, Schulze KL, Zhai RG, Hiesinger PR, Zhou Y, Mehta SQ, Cao Y, Roos J, Bellen HJ
Neuron (2003) 40(4): 733-48.
- Endophilin and synaptojanin hook up to promote synaptic vesicle endocytosis.
Song W, Zinsmaier KE
Neuron (2003) 40(4): 665-7.
- Synaptojanin 1 contributes to maintaining the stability of GABAergic transmission in primary cultures of cortical neurons.
Luthi A, Di Paolo G, Cremona O, Daniell L, De Camilli P, McCormick DA
The Journal of neuroscience : the official journal of the Society for Neuroscience (2001) 21(23): 9101-11.
- Mutations in synaptojanin disrupt synaptic vesicle recycling.
Harris TW, Hartweg E, Horvitz HR, Jorgensen EM
The Journal of cell biology (2000) 150(3): 589-600.
- Direct interaction of the 170 kDa isoform of synaptojanin 1 with clathrin and with the clathrin adaptor AP-2.
Haffner C, Di Paolo G, Rosenthal JA, de Camilli P
Current biology : CB (2000) 10(8): 471-4.
- Synaptojanin is the major constitutively active phosphatidylinositol-3,4,5-trisphosphate 5-phosphatase in rodent brain.
Woscholski R, Finan PM, Radley E, Totty NF, Sterling AE, Hsuan JJ, Waterfield MD, Parker PJ
The Journal of biological chemistry (1997) 272(15): 9625-8.
- Identification of the major synaptojanin-binding proteins in brain.
de Heuvel E, Bell AW, Ramjaun AR, Wong K, Sossin WS, McPherson PS
The Journal of biological chemistry (1997) 272(13): 8710-6.
- Tissue-specific alternative splicing generates two synaptojanin isoforms with differential membrane binding properties.
Ramjaun AR, McPherson PS
The Journal of biological chemistry (1996) 271(40): 24856-61.